arXiv:1201.0190v1 [math.DG] 30 Dec 2011

TRANSFORMATIONS OF HARMONIC BUNDLES AND WILLMORE SURFACES

A. C. QUINTINO

Abstract. Willmore surfaces are the extremals of the Willmore functional (possibly under a constraint on the conformal structure). With the characterization of Willmore surfaces by the harmonicity of the mean curvature sphere congruence ([13], [18]), a zero-curvature formulation follows ([5]). Deformations on the level of bundles prove to give rise to deformations on the level of surfaces, with the definition of a spectral deformation ([5]) and of a Bäcklund transformation ([9]) of Willmore surfaces into new ones, with a permutability between the two ([9]). This note is dedicated to a self-contained account of the topic in the light-cone picture.

1. Introduction

Among the classes of Riemannian submanifolds, there is that of Willmore surfaces, named after T. Willmore [22] (1965), although the topic was mentioned by W. Blaschke [1] (1929) and by G. Thomsen [20] (1923), as a variational problem of optimal realization of a given surface in 3-space. Early in the nineteenth century, S. Germain [14], [15] studied elastic surfaces. On her pioneering analysis, she claimed that the elastic force of a thin plate is proportional to its mean curvature. Since then, the mean curvature remains a key concept in theory of elasticity. In modern literature on the elasticity of membranes, a weighted sum of the total mean curvature, the total squared mean curvature and the total Gaussian curvature is considered the bending energy of a membrane. By neglecting the total mean curvature, by physical considerations, and having in consideration Gauss-Bonnet Theorem, T. Willmore defined the Willmore [bending] energy of a compact oriented surface Σ, without boundary, immersed in $\mathbb{R}^3$ to be $\int_\Sigma H^2 dA$, averaging the mean curvature square over the surface. The Willmore energy "extends" to surfaces immersed in a general Riemannian manifold $\bar{M}$ of constant sectional curvature by means of

$$\mathcal{W} := \int_\Sigma |\Pi^0|^2 dA,$$

the total squared norm of the trace-free part Π^0 of the second fundamental form. In fact, given $(X_i)_{i=1,2}$ a local orthonormal frame of $T\Sigma$, Gauss equation establishes

$$|\Pi^0|^2 := \sum_{i,j} (\Pi^0(X_i, X_j), \Pi^0(X_i, X_j)) = 2(|\mathcal{H}|^2 - K + \bar{K}),$$

for $\mathcal{H}$ the mean curvature vector, K the Gaussian curvature of Σ and $\bar{K}$ the sectional curvature of $\bar{M}$. In particular, for surfaces in $\mathbb{R}^3$, the functional defined by $\mathcal{W}$ shares critical points with the one defined by the total squared mean curvature. Willmore surfaces are the extremals of the Willmore functional. Constrained Willmore surfaces appear as the generalization of Willmore surfaces that arises when we consider extremals of the Willmore functional with respect to infinitesimally conformal variations, those that infinitesimally preserve the conformal structure, that is,

$$\frac{d}{dt}_{|t=0} (Z^{1,0}, Z^{1,0})_t = 0,$$

for the variation $(,)_t$ of the induced metric (or rather its complex bilinear extension), fixing $Z^{1,0}$ a $(1,0)$-vector field; rather than with respect to all variations.

A classical result by Thomsen [20] characterizes isothermic Willmore surfaces in 3-space as minimal surfaces in some 3-dimensional space-form. Constant mean curvature surfaces in 3-dimensional space-forms are examples of isothermic constrained Willmore surfaces, as proven by J. Richter [17]. However, isothermic constrained Willmore surfaces in 3-space are not necessarily constant mean curvature surfaces in some space-form, as established by an example, presented in [2], of a constrained Willmore cylinder that does not have constant mean curvature in any space-form.

Under a conformal change $e^{2u}g$ of a metric g, for some function u, the second fundamental form Π of an isometric immersion f changes via

$$\Pi_{e^{2u}g}(X, Y) = \Pi_g(X, Y) - g(X, Y)\pi_N(f^*(du)^*), \tag{1}$$

for π_N the normal projection and $f^*(du)^*$ the pull-back by f of the contravariant form $(du)^*$ form of du with respect to g (see, for example, [23], section 3.12). Hence, under a conformal change of the metric, the trace-free part of the second fundamental form remains invariant, so that its squared norm and the area element change in an inverse way, leaving the Willmore energy unchanged. In particular, this establishes the class of (constrained) Willmore surfaces as a Möbius invariant class.

A very well known result ([1] ($n = 3$), [13], [18] (general n)) characterizes Willmore surfaces by the harmonicity of the central sphere congruence. With this characterization, a zero-curvature formulation of Willmore surfaces follows, according to the zero-curvature formulation of harmonicity of a map into a Grassmanian by Uhlenbeck [21]. This characterization generalizes [5] to constrained Willmore surfaces and deformations on the level of harmonic bundles prove to give rise to deformations on the level of surfaces. The class of constrained Willmore surfaces is established as a class of surfaces with strong links to the theory of integrable systems, admitting a spectral deformation, defined by the action of a loop of flat metric connections, as in [5], and Bäcklund transformations, defined by applying a dressing action, as studied in [9]. A permutability between the two is established in [9]. All these transformations corresponding to the zero Lagrange multiplier preserve the class of Willmore surfaces. The isothermic surface condition is known to be preserved under constrained Willmore spectral deformation, cf. [8]. As for Bäcklund transformation of constrained Willmore surfaces, we believe it does not necessarily preserve the isothermic condition. In contrast, the constancy of the mean curvature of a surface in 3-dimensional space-form is preserved by both constrained Willmore spectral deformation and Bäcklund transformation, for special choices of parameters, with preservation of both the space-form and the mean curvature in the latter case, cf. [16].

This note is dedicated to a self-contained account of the topic in the light-cone picture. The research work was developed as part of the Ph.D. studies of the author at the University of Bath, UK, under the supervision of Prof. F. E. Burstall.

2. Constrained Willmore surfaces and constrained harmonicity

Consider $\underline{\mathbb{C}}^{n+2} = \Sigma \times (\mathbb{R}^{n+1,1})^{\mathbb{C}}$ provided with the complex bilinear extension of the metric on $\underline{\mathbb{R}}^{n+1,1}$. In what follows, we shall make no explicit distinction between a bundle and its complexification, and move from real tensors to complex tensors by complex multilinear extension, with no need for further reference, preserving notation.

Throughout this text, we will consider the identification $\wedge^2\mathbb{R}^{n+1,1} \cong o(\mathbb{R}^{n+1,1})$ of the exterior power $\wedge^2\mathbb{R}^{n+1,1}$ with the orthogonal algebra $o(\mathbb{R}^{n+1,1})$ via $u\wedge v(w) := (u,w)v - (v,w)u$ for $u, v, w \in \mathbb{R}^{n+1,1}$. We consider the bundle $\mathrm{End}(\underline{\mathbb{R}}^{n+1,1})$ - and, more generally, any bundle of morphisms - provided with the metric defined by $(\xi, \eta) := \mathrm{tr}(\eta^t\xi)$ and we shall move from a connection on $\underline{\mathbb{R}}^{n+1,1}$ to a connection on $\mathrm{End}(\underline{\mathbb{R}}^{n+1,1})$ via $\nabla\xi = \nabla \circ \xi - \xi \circ \nabla$, with preservation of notation. Note that, in the case of a metric connection ∇ on $\underline{\mathbb{R}}^{n+1,1}$, we have $\nabla(u \wedge v) = \nabla u \wedge v + u \wedge \nabla v$, for all $u, v \in \Gamma(\underline{\mathbb{R}}^{n+1,1})$.

Our theory is local and, throughout this text, with no need for further reference, restriction to a suitable non-empty open set shall be underlying.

2.1. **Real constrained Willmore surfaces.**

2.1.1. *Conformal geometry of the sphere.* Our study is one of surfaces in n-dimensional space-forms, with $n \geq 3$, from a conformally-invariant viewpoint. For this, we find a convenient setting in Darboux's light-cone model of the conformal n-sphere [12], viewing the n-sphere not as the round sphere in the Euclidean space $\mathbb{R}^{n+1}$ but as the celestial sphere in the Lorentzian spacetime $\mathbb{R}^{n+1,1}$. So contemplate the light-cone $\mathcal{L}$ in the Lorentzian vector space $\mathbb{R}^{n+1,1}$ and its projectivisation $\mathbb{P}(\mathcal{L})$, provided with the conformal structure defined by a metric g_σ arising from a never-zero section σ of the tautological bundle $\pi : \mathcal{L} \to \mathbb{P}(\mathcal{L})$ via

$$g_\sigma(X, Y) = (d\sigma(X), d\sigma(Y)).$$

For $v_\infty \in \mathbb{R}^{n+1,1}_\times$, set $S_{v_\infty} := \{v \in \mathcal{L} : (v, v_\infty) = -1\}$, an n-dimensional submanifold $\mathbb{R}^{n+1,1}$ which inherits from $\mathbb{R}^{n+1,1}$ a positive definite metric of (constant) sectional curvature $-(v_\infty, v_\infty)$. In fact, for v_∞ non-null, orthoprojection onto $\langle v_\infty\rangle^\perp$ induces an isometry between S_{v_∞} and $\{v \in \langle v_\infty\rangle^\perp : (v, v) = -1/(v_\infty, v_\infty)\}$, whereas, when v_∞ is null, for any choice of $v_0 \in S_{v_\infty}$, orthoprojection onto $\langle v_0, v_\infty\rangle^\perp$ restricts to an isometry of S_{v_∞}. By construction, the bundle projection π restricts to give a conformal diffeomorphism $\pi_{|S_{v_\infty}} : S_{v_\infty} \to \mathbb{P}(\mathcal{L})\backslash\mathbb{P}(\mathcal{L} \cap \langle v_\infty\rangle^\perp)$. In particular, choosing v_∞ time-like identifies $\mathbb{P}(\mathcal{L})$ with the conformal n-sphere. The beauty of this model is that it *linearises* the conformal geometry of the sphere. For example, k-spheres in $S^n \cong \mathbb{P}(\mathcal{L})$ are identified with $(k + 1, 1)$-planes V in $\mathbb{R}^{n+1,1}$ via $V \mapsto \mathbb{P}(\mathcal{L} \cap V) \subset \mathbb{P}(\mathcal{L})$.

2.1.2. *Surfaces in the light-cone picture.* For us, a mapping $\Lambda : \Sigma \to S^n \cong \mathbb{P}(\mathcal{L})$, of a surface Σ, is the same as a null line subbundle of the trivial bundle $\underline{\mathbb{R}}^{n+1,1} = \Sigma \times \mathbb{R}^{n+1,1}$, in the natural way. Given such a Λ, we define

$$\Lambda^{(1)} := \langle \sigma, d\sigma(T\Sigma)\rangle,$$

for σ a lift of Λ. Note that Λ is an immersion if and only if the bundle $\Lambda^{(1)}$ has rank 3.

Let then $\Lambda : \Sigma \to S^n \cong \mathbb{P}(\mathcal{L})$ be an immersion of an oriented surface Σ, which we provide with the conformal structure $\mathcal{C}_\Lambda$ induced by Λ and with J the canonical complex structure (that is, 90° rotation in the positive direction in the tangent spaces, a notion that is obviously invariant under conformal changes of the metric). Note that every lift $\sigma : \Sigma \to \mathbb{R}^{n+1,1}$ of Λ is conformal. Set

$$\Lambda^{1,0} := \Lambda \oplus d\sigma(T^{1,0}\Sigma), \quad \Lambda^{0,1} := \Lambda \oplus d\sigma(T^{0,1}\Sigma), \tag{2}$$

independently of the choice of a lift σ of Λ, defining in this way two complex rank 2 subbundles of $\Lambda^{(1)}$, complex conjugate of each other. The nullity and conformality of the lifts of Λ establish the isotropy of the bundles $\Lambda^{1,0}$ and $\Lambda^{0,1}$, whilst the fact that Λ is an immersion establishes that $\Lambda^{1,0}$ and $\Lambda^{0,1}$ intersect in Λ.

We restrict our study to surfaces in S^n which are not contained in any subsphere of S^n. This ensures, in particular, that, given $v_\infty \in \mathbb{R}^{n+1,1}$ non-zero, $\Lambda(\Sigma) \nsubseteq \mathbb{P}(\mathcal{L} \cap \langle v_\infty\rangle^\perp)$, and we define a local immersion

$$\sigma_\infty := \pi_{|S_{v_\infty}} \circ \Lambda = \frac{-1}{(\sigma, v_\infty)}\,\sigma : \Sigma \to S_{v_\infty},$$

into the space-form S_{v_∞}, independently of σ a lift of Λ.

A fundamental construction in conformal geometry of surfaces is the mean curvature sphere congruence, the bundle of 2-spheres tangent to the surface and sharing mean curvature vector with it at each point (although the mean curvature vector is not conformally-invariant, under a conformal change of the metric it changes in the same way for the surface and the osculating 2-sphere). From the early twentieth century, the family of the mean curvature spheres of a surface was known as the central sphere congruence, cf. W. Blaschke [1]. Nowadays, after R. Bryant's paper [3], it goes as well by the name conformal Gauss map, although the central sphere congruence carries, not only first order contact information, but second order as well.

Let $S : \Sigma \to \mathcal{G} := \mathrm{Gr}_{(3,1)}(\mathbb{R}^{n+1,1})$ be the central sphere congruence of Λ,

$$S = \Lambda^{(1)} \oplus \langle \triangle\sigma \rangle = \langle \sigma, \sigma_z, \sigma_{\bar{z}}, \sigma_{z\bar{z}} \rangle,$$

for σ a lift of Λ, $\triangle\sigma$ the Laplacian of σ with respect to the metric g_σ and z a holomorphic chart of Σ. Let π_S and $\pi_{S^\perp}$ denote the orthogonal projections of $\underline{\mathbb{R}}^{n+1,1}$ onto S and $S^\perp$, respectively. Given z a holomorphic chart of Σ, let g_z denote the metric induced in Σ by z. Note that, by the conformality of z, $(\sigma, \sigma_{z\bar{z}}) = -(\sigma_z, \sigma_{\bar{z}})$ is never-zero. In many occasions, it will be useful to consider a special choice of lift of Λ, the *normalized* lift with respect to z, the section $\sigma^z : \Sigma \to \mathcal{L}^+$ of Λ (given a choice $\mathcal{L}^+$ of one of the two connected components of $\mathcal{L}$) defined by $g_{\sigma^z} = g_z$. For further reference, note that this condition establishes, in particular, that $(\sigma^z_z, \sigma^z_{\bar{z}})$ is constant, $(\sigma^z_z, \sigma^z_{\bar{z}}) = \frac{1}{2}$, and, therefore,

$$\pi_S \sigma^z_{zz} \in \Gamma((\Lambda^{(1)})^\perp \cap S) = \Gamma\Lambda. \tag{3}$$

Consider the decomposition of the trivial flat connection d on $\underline{\mathbb{R}}^{n+1,1}$ as

$$d = \mathcal{D} \oplus \mathcal{N}$$

for $\mathcal{D}$ the connection given by the sum of the connections ∇^S and $\nabla^{S^\perp}$ induced on S and $S^\perp$, respectively, by d. Note that $\mathcal{D}$ is a metric connection and, therefore, $\mathcal{N}$ is skew-symmetric, $\mathcal{N} \in \Omega^1(S \wedge S^\perp)$. Note that, given $\xi \in \Gamma(S \wedge S^\perp)$, the transpose of $\xi|_S$ is $-\xi|_{S^\perp}$, and define a bundle isomorphism $S \wedge S^\perp \to \mathrm{Hom}(S, S^\perp)$ by $\eta \mapsto \eta|_S$. Together with the canonical identification of $S^*T\mathcal{G}$ and $\mathrm{Hom}(S, S^\perp)$, via $X \mapsto (\rho \mapsto \pi_{S^\perp}(d_X\rho))$, this gives an identification

$$S^*T\mathcal{G} \cong \mathrm{Hom}(S, S^\perp) \cong S \wedge S^\perp, \tag{4}$$

of bundles provided with the canonical metrics and connections (for the connection $\mathcal{D}$ on $\underline{\mathbb{R}}^{n+1,1}$), (see, for example, [10]), which we will consider throughout. Observe that, under the identification (4),

$$dS = \mathcal{N}. \tag{5}$$

2.1.3. *Willmore surfaces and harmonicity.* Suppose, for the moment, that Σ is compact. The next result follows the definition presented in [7], in the quaternionic setting, for the particular case of $n = 4$. A proof is presented below.

Theorem 1. *[[7]]*

$$\mathcal{W}(\Lambda)=\frac{1}{2}\int_{\Sigma}(dS\wedge *dS), \tag{6}$$

for the 2*-form* $(\wedge)$ *defined from the metric on* $S^*T\mathcal{G}$*.*

Proof. By (5), fixing a metric on Σ, $|dS|^2=|\mathcal{N}|^2$. To prove the theorem, we fix $v_\infty\in\mathbb{R}^{n+1,1}$ non-zero, provide Σ with the metric induced by σ_∞ and show that $|\mathcal{N}|^2=2|\Pi^0_\infty|^2$, for Π^0_∞ the trace-free part of the second fundamental form of σ_∞.

Fixing a local orthonormal frame $\{X_i\}_i$ of $T\Sigma$, we have

$$|\mathcal{N}|^2=-\sum_i\operatorname{tr}(\mathcal{N}_{X_i}\mathcal{N}_{X_i})=2\sum_i\operatorname{tr}(\mathcal{N}^t_{X_i}\mathcal{N}_{X_i}|_S).$$

Recall that if $(e_i)_i$ and $(\hat{e}_i)_i$ are dual basis of a vector space E provided with a metric $(,)$, then, given $\mu\in\operatorname{End}(E)$, $\operatorname{tr}(\mu)=\sum_i(\mu(e_i),\hat{e}_i)$. Let $\hat{\sigma}_\infty$ be the section of S determined by the conditions $(\hat{\sigma}_\infty,\hat{\sigma}_\infty)=0$, $(\sigma_\infty,\hat{\sigma}_\infty)=-1$ and $(\hat{\sigma}_\infty,d\sigma_\infty)=0$. Then $(\sigma_\infty,d_{X_1}\sigma_\infty,d_{X_2}\sigma_\infty,\hat{\sigma}_\infty)$ is a frame of S with dual $(-\hat{\sigma}_\infty,d_{X_1}\sigma_\infty,d_{X_2}\sigma_\infty,-\sigma_\infty)$ and we conclude that

$$|\mathcal{N}|^2=2\sum_{i,j}(\mathcal{N}_{X_i}(d_{X_j}\sigma_\infty),\mathcal{N}_{X_i}(d_{X_j}\sigma_\infty)).$$

The bundle N_∞ normal to the surface σ_∞ can be identified with the normal bundle to the central sphere congruence of Λ via the isomorphism $\mathcal{Q}_\infty : N_\infty\rightarrow S^\perp$, of bundles provided with a metric and a connection, defined by $\xi\mapsto\xi+(\xi,\mathcal{H}_\infty)\sigma_\infty$, for $\mathcal{H}_\infty$ the mean curvature vector of σ_∞, cf. [8]. For $\xi\in N_\infty\subset\sigma^*_\infty TS_{v_\infty}=\langle\sigma_\infty,v_\infty\rangle^\perp$,

$$(\mathcal{Q}_\infty(\xi),\pi_{S^\perp}(v_\infty))=(\xi,v_\infty)+(\xi,\mathcal{H}_\infty)(\sigma_\infty,v_\infty)=-(\mathcal{Q}_\infty(\xi),\mathcal{Q}_\infty(\mathcal{H}_\infty)),$$

establishing $\pi_{S^\perp}(v_\infty)=-\mathcal{Q}_\infty(\mathcal{H}_\infty)$. Hence $\mathcal{N}_{X_i}(d_{X_j}\sigma_\infty)=\mathcal{Q}_\infty(\Pi^0_\infty(X_i,X_j)$, completing the proof. □

The intervention of the conformal structure in (6) restricts to the Hodge $*$-operator, which is conformally-invariant on 1-forms over a surface. Theorem 1 presents a manifestly conformally-invariant formulation of the Willmore energy.

Theorem 1 makes it clear that

$$\mathcal{W}(\Lambda)=E(S,\mathcal{C}_\Lambda),$$

the Willmore energy of Λ coincides with the Dirichlet energy of S with respect to any of the metrics in the conformal class $\mathcal{C}_\Lambda$ (although the Levi-Civita connection is not conformally-invariant, the Dirichlet energy of a mapping of a surface is preserved under conformal changes of the metric (and so is its harmonicity)). Furthermore, in a very well known result:

Theorem 2 ([1] ($n=3$), [13], [18] (general n))**.** Λ *is a Willmore surface if and only if its central sphere congruence* $S:(\Sigma,\mathcal{C}_\Lambda)\rightarrow\mathcal{G}$ *is a harmonic map.*

Next we present a proof of Theorem 2 in the light-cone picture. This is a generalization of the proof presented in [7], in the quaternionic setting, for the particular case of $n = 4$.

Proof. Given a variation Λ_t of Λ and S_t the corresponding variation of S through central sphere congruences, the Dirichlet energy $E(S_t, \mathcal{C}_t)$ of S_t with respect to the conformal structure $\mathcal{C}_t$ induced in Σ by Λ_t is given by $\frac{1}{2}\int_M (dS_t \wedge *_t dS_t)$, for dA_t and $*_t$ the area element and the Hodge $*$-operator of (Σ, g_t), respectively, fixing $g_t \in \mathcal{C}_t$. Hence

$$\frac{d}{dt}_{|t=0} E(S_t, \mathcal{C}_t) = \frac{1}{2}\int_M ((d\dot{S} \wedge *dS) + (dS \wedge \dot{*}dS) + (dS \wedge *d\dot{S})),$$

abbreviating $\frac{d}{dt}_{|t=0}$ by a dot. Let $(J_t)_t$ be the corresponding variation of J through canonical complex structures. Differentiation at $t = 0$ of $*_t dS_t = -(dS_t)J_t$ gives $\dot{*}dS = -(dS)\dot{J}$, whilst that of $J_t^2 = -I$ gives $\dot{J}J = -J\dot{J}$ and, in particular, that $\dot{J}$ intertwines the eigenspaces of J. The $\mathcal{C}_\Lambda$-conformality of S, $(d_{X \pm iJX}S, d_{X \pm iJX}S) = 0$, respectively, for $X \in \Gamma(TM)$, establishes then $(dS \wedge \dot{*}dS) = 0$ and, therefore,

$$\frac{d}{dt}_{|t=0} E(S_t, \mathcal{C}_t) = \frac{d}{dt}_{|t=0} E(S_t, \mathcal{C}_0).$$

It is now clear that if $S : (\Sigma, \mathcal{C}_\Lambda) \to \mathcal{G}$ is harmonic then Λ is Willmore.

Conversely, suppose Λ is Willmore, fix z a holomorphic chart of $(\Sigma, \mathcal{C}_\Lambda)$ and let us show that the tension field τ_z of $S : (\Sigma, g_z) \to \mathcal{G}$ vanishes. First of all, observe that $4\nabla_{\delta_z} S_{\bar{z}} = \tau_z = 4\nabla_{\delta_{\bar{z}}} S_z$ to conclude that $\Lambda^{(1)} \subset \ker \tau_z$: by (3), $(\nabla_{\delta_z} S_{\bar{z}})\sigma^z_z = \nabla^{S^\perp}_{\delta_z}(\pi_{S^\perp}\sigma^z_{z\bar{z}}) - \pi_{S^\perp}(\nabla^S_{\delta_z}\sigma^z_z)_{\bar{z}} = 0$, and, similarly, $(\nabla_{\delta_{\bar{z}}} S_z)\sigma^z_{\bar{z}} = 0$, whilst $(\nabla_{\delta_z} S_{\bar{z}})\sigma^z = 0$ is immediate. It follows that $\mathrm{Im}\tau^t_z \subset \Lambda$. Fix $\Lambda_t = \langle \sigma_t \rangle$ a variation of Λ and let S_t be the corresponding variation of S through central sphere congruences. Then $\tau^t_z(\pi_{S^\perp}\dot{\sigma}) = \lambda\sigma_0$, for some $\lambda \in C^\infty(\Sigma, \mathbb{R})$, and, therefore, $\mathrm{tr}(\tau^t_z \dot{S}) = \lambda$ (note that $\dot{S}\sigma = \pi_{S^\perp}\dot{\sigma}$). On the other hand, classically, $0 = \frac{d}{dt}_{|t=0} E(S_t, \mathcal{C}_\Lambda) = -\int_\Sigma (\dot{S}, \tau_z) dA_z = -\int_\Sigma \mathrm{tr}(\tau^t_z \dot{S}) dA_z$, for dA_z the area element of (Σ, g_z). Now suppose τ_z is non-zero. Then so is $\tau^t_z \in \Gamma(\mathrm{Hom}(S^\perp, S))$, so we can choose σ_t such that λ is positive, which leads to a contradiction and completes the proof. □

Having characterized Willmore surfaces by the harmonicity of the central sphere congruence, and recalling (5), we deduce the Willmore surface equation,

$$d^{\mathcal{D}} * \mathcal{N} = 0. \tag{7}$$

More generally, we have a manifestly conformally-invariant characterization of constrained Willmore surfaces in space-forms, first established in [8] and reformulated in [5] as follows:

Theorem 3 ([5])**.** *Λ is a constrained Willmore surface if and only if there exists a real form $q \in \Omega^1(\Lambda \wedge \Lambda^{(1)})$ with*

$$d^{\mathcal{D}} q = 0 \tag{8}$$

such that

$$d^{\mathcal{D}} * \mathcal{N} = 2\,[q \wedge *\mathcal{N}]. \tag{9}$$

We refer to such a form q as a [Lagrange] multiplier *for Λ and say that Λ is a q-*constrained Willmore surface*.*

A proof of Theorem 3 can be found in [5].

Sketch proof Calculus of variations techniques show that the variational Willmore energy relates to the variational surface by $\dot{\mathcal{W}} = \int_\Sigma ((d^{\mathcal{D}} * \mathcal{N}) \wedge \dot{\Lambda})$, for some non-degenerate pairing $(\,\wedge\,)$. For general variations, $\dot{\Lambda}$ can be arbitrary, establishing the Willmore surface equation (7), whereas infinitesimally conformal variations are characterized by the normal variational being in the image of the conformal Killing operator $\overline{\bar{\partial}}$, which, according to Weyl's Lemma, consists of $(H^0K)^{\perp}$, for H^0K the space of holomorphic quadratic differentials. The result follows by defining a multiplier q from a quadratic differential $q^z dz^2 \in (H^0K)^{\perp}$ via $q_{\delta_z}\sigma_z = -\frac{1}{2}\,q^z\sigma$, for σ a lift of Λ and z a holomorphic chart of Σ.

Willmore surfaces are the 0-constrained Willmore surfaces. The zero multiplier is not necessarily the only multiplier for a constrained Willmore surface with no constraint on the conformal structure, though. In fact, the uniqueness of multiplier characterizes non-isothermic constrained Willmore surfaces, as we shall see below in this text.

The characterization of constrained Willmore surfaces above motivates a natural extension to surfaces that are not necessarily compact.

Next we present a useful result, which establishes, in particular, that if q is a multiplier for Λ, then $q^{1,0}$ takes values in $\Lambda \wedge \Lambda^{0,1}$.

Proposition 4. *Given $q \in \Omega^1(\Lambda \wedge \Lambda^{(1)})$ real,*

i) if $d^{\mathcal{D}}q = 0$ then $q^{1,0} \in \Omega^{1,0}(\Lambda \wedge \Lambda^{0,1})$ or, equivalently, $q^{0,1} \in \Omega^{0,1}(\Lambda \wedge \Lambda^{1,0})$;

ii) $d^{\mathcal{D}}q = 0$ if and only if $d^{\mathcal{D}}q^{1,0} = 0$, or, equivalently, $d^{\mathcal{D}}q^{0,1} = 0$;

iii) $d^{\mathcal{D}}q = 0$ if and only if $d^{\mathcal{D}} * q = 0$.

Proof. Fix z a holomorphic chart of Σ. First of all, observe that a section ξ of $\Lambda \wedge \Lambda^{(1)}$ is a section of $\Lambda \wedge \Lambda^{1,0}$ if and only if $\xi(\sigma^z_{\bar{z}}) = 0$. Suppose $d^{\mathcal{D}}q = 0$. Then, in particular, $d^{\mathcal{D}}q\,(\delta_z, \delta_{\bar{z}})\,\sigma^z = 0$, or, equivalently, $\mathcal{D}_{\delta_z}(q_{\delta_{\bar{z}}}\sigma^z) - q_{\delta_{\bar{z}}}(\mathcal{D}_{\delta_z}\sigma^z) - \mathcal{D}_{\delta_{\bar{z}}}(q_{\delta_z}\sigma^z) + q_{\delta_z}(\mathcal{D}_{\delta_{\bar{z}}}\sigma^z) = 0$, establishing

$$q_{\delta_z}\sigma^z_{\bar{z}} = q_{\delta_{\bar{z}}}\sigma^z_z. \tag{10}$$

In its turn, $d^{\mathcal{D}}q\,(\delta_z, \delta_{\bar{z}})\,\sigma^z_z = 0$ implies $\mathcal{D}_{\delta_z}(q_{\delta_{\bar{z}}}\sigma^z_z) - \mathcal{D}_{\delta_{\bar{z}}}(q_{\delta_z}\sigma^z_z) + q_{\delta_z}\sigma^z_{z\bar{z}} = 0$, by (3). On the other hand, the skew-symmetry of q establishes $(q\sigma^z_{z\bar{z}}, \sigma^z_{z\bar{z}}) = 0$ and, therefore,

$$q\sigma^z_{z\bar{z}} = \mu\sigma_z + \eta\sigma^z_{\bar{z}}, \tag{11}$$

for some $\mu, \eta \in \Omega^1(\underline{\mathbb{C}})$. Hence $\mathcal{D}_{\delta_z}(q_{\delta_{\bar z}}\sigma^z_z) + \mu_{\delta_z}\sigma^z_z = \mathcal{D}_{\delta_{\bar z}}(q_{\delta_z}\sigma^z_z) - \eta_{\delta_z}\sigma^z_{\bar z}$. It is obvious that a section of $\Lambda \wedge \Lambda^{(1)}$ transforms sections of $\Lambda^{(1)}$ into sections of Λ, so that, in particular, both $q_{\delta_{\bar z}}\sigma^z_z$ and $q_{\delta_z}\sigma^z_z$ are sections of Λ. We conclude that $\mathcal{D}_{\delta_z}(q_{\delta_{\bar z}}\sigma^z_z) + \mu_{\delta_z}\sigma^z_z$ is a section of $\Lambda^{1,0} \cap \Lambda^{0,1} = \Lambda$. Write $q_{\delta_{\bar z}}\sigma^z_z = \lambda\sigma^z$, with $\lambda \in \Gamma(\underline{\mathbb{C}})$. Then $\lambda_z\sigma^z + (\lambda + \mu_{\delta_z})\sigma^z_z = \gamma\sigma^z$, for some $\gamma \in \Gamma(\underline{\mathbb{C}})$. In particular, $\lambda = -\mu_{\delta_z}$. Equation (11) establishes, on the other hand, $q = -2\mu\,\sigma^z \wedge \sigma^z_z - 2\eta\,\sigma^z \wedge \sigma^z_{\bar z}$ and, in particular, $q_{\delta_z}\sigma^z_{\bar z} = \mu_{\delta_z}\sigma^z$. Equation (10) completes the proof of *i*).

Next we prove *ii*). By (3), $\mathcal{D}^{1,0}\Gamma(\Lambda^{1,0}) \subset \Omega^{1,0}(\Lambda^{1,0})$ or, equivalently, $\mathcal{D}^{0,1}\Gamma(\Lambda^{0,1}) \subset \Omega^{0,1}(\Lambda^{0,1})$ and, therefore, following *i*), $d^{\mathcal{D}}q^{1,0} \in \Omega^2(\Lambda \wedge \Lambda^{0,1})$ and $d^{\mathcal{D}}q^{0,1} \in \Omega^2(\Lambda\wedge\Lambda^{1,0})$. Hence $d^{\mathcal{D}}q = 0$ forces $d^{\mathcal{D}}q^{1,0}$ and $d^{\mathcal{D}}q^{0,1}$ to vanish separately. The reality of q completes the proof of *ii*).

As for *iii*), it is immediate from *ii*). □

With a characterization of Willmore surfaces by the harmonicity of the central sphere congruence, a zero curvature formulation follows, according to the zero-curvature formulation of harmonicity of a map into a Grassmanian by Uhlenbeck [21]. More generally:

Theorem 5 ([5]). *Λ is a constrained Willmore surface if and only if there exists $q \in \Omega^1(\Lambda \wedge \Lambda^{(1)})$ such that*

$$d^\lambda_q := \mathcal{D} + \lambda^{-1}\mathcal{N}^{1,0} + \lambda\mathcal{N}^{0,1} + (\lambda^{-2} - 1)q^{1,0} + (\lambda^2 - 1)q^{0,1}$$

is flat for all $\lambda \in S^1$. Such a form q is said to be a [Lagrange] multiplier *for Λ and Λ is said to be a q-*constrained Willmore surface*. Willmore surfaces are the 0-constrained Willmore surfaces.*

A proof of Theorem 5 can be found in [5].

Sketch proof According to the decomposition

$$o(\underline{\mathbb{R}}^{n+1,1}) = (\wedge^2 S \oplus \wedge^2 S^\perp) \oplus S \wedge S^\perp, \tag{12}$$

the flatness of d, characterized by $0 = R^{\mathcal{D}} + d^{\mathcal{D}}\mathcal{N} + \frac{1}{2}\,[\mathcal{N} \wedge \mathcal{N}]$, for $[\wedge]$ the 2-form defined from the Lie Bracket in $o(\mathbb{R}^{n+1,1})$, encodes two structure equations, namely, $R^{\mathcal{D}} + \frac{1}{2}\,[\mathcal{N} \wedge \mathcal{N}] = 0$ and $d^{\mathcal{D}}\mathcal{N} = 0$. The proof follows by computing the curvature of the connections.

2.2. Complexified constrained Willmore surfaces.

The transformations of a constrained Willmore surface Λ we present below in this work are, in particular, pairs $((\Lambda^{1,0})^*, (\Lambda^{0,1})^*)$ of transformations $(\Lambda^{1,0})^*$ and $(\Lambda^{0,1})^*$ of $\Lambda^{1,0}$ and $\Lambda^{0,1}$, respectively. The fact that $\Lambda^{1,0}$ and $\Lambda^{0,1}$ intersect in a rank 1 bundle will ensure that $(\Lambda^{1,0})^*$ and $(\Lambda^{0,1})^*$ have the same property. The isotropy of $\Lambda^{1,0}$ and $\Lambda^{0,1}$ will ensure that of $(\Lambda^{1,0})^*$ and $(\Lambda^{0,1})^*$ and, therefore, that of their intersection. The reality of the bundle $\Lambda^{1,0} \cap \Lambda^{0,1}$ is preserved by the spectral deformation, but it is not clear that the same is necessarily true for Bäcklund transformation. This motivates the definition of *complexified surface*.

Fix a conformal structure $\mathcal{C}$ on Σ and consider the corresponding complex structure on Σ. Let $\hat{d}$ be a flat metric connection on $\underline{\mathbb{C}}^{n+2}$ and d denote the trivial flat connection. In what follows, omitting the reference to some specific connection shall be understood as an implicit reference to d.

Definition 1. *We define a* complexified $\hat{d}$-surface *to be a pair* $(\Lambda^{1,0},\Lambda^{0,1})$ *of isotropic rank* 2 *subbundles of* $\underline{\mathbb{C}}^{n+2}$ *intersecting in a rank* 1 *bundle* $\Lambda := \Lambda^{1,0}\cap\Lambda^{0,1}$ *such that*

$$\hat{d}^{1,0}\Gamma\Lambda \subset \Omega^{1,0}\Lambda^{1,0}, \quad \hat{d}^{0,1}\Gamma\Lambda \subset \Omega^{0,1}\Lambda^{0,1}.$$

Obviously, given Λ a (real) surface in $\mathbb{P}(\mathcal{L})$, $(\Lambda^{1,0},\Lambda^{0,1})$ is a complexified surface with respect to $\mathcal{C}_\Lambda$. Henceforth, we drop the term "complexified" and use *real surface* when referring explicitly to a complexified surface $(\Lambda^{1,0},\Lambda^{0,1})$ defining a real surface Λ. Observe that $(\Lambda^{1,0},\Lambda^{0,1})$ is a real surface if and only if Λ is a real bundle (recall that Λ is an immersion if and only if the bundle $\Lambda^{1,0}+\Lambda^{0,1}$ has rank 3).

Observe, on the other hand, that, in the particular case of a real surface $(\Lambda^{1,0},\Lambda^{0,1})$, the notation $\Lambda^{1,0}$ and $\Lambda^{0,1}$ is consistent with (2). Indeed, the $\mathcal{C}$-isotropy of $\Lambda^{1,0}$ characterizes the $\mathcal{C}$-conformality of the lifts of Λ, or equivalently, the fact that $\mathcal{C}=\mathcal{C}_\Lambda$.

2.2.1. *Constrained Willmore surfaces and constrained harmonicity.* Theorem 5 motivates the notion of *constrained harmonicity* of a bundle, which we present next and which will apply to the central sphere congruence to provide a zero-curvature formulation of constrained Willmore surfaces.

Given V a non-degenerate subbundle of $\underline{\mathbb{C}}^{n+2}$, consider the decomposition $\hat{d} = \mathcal{D}_V^{\hat{d}} + \mathcal{N}_V^{\hat{d}}$ for $\mathcal{D}_V^{\hat{d}}$ the metric connection on $\underline{\mathbb{C}}^{n+2}$ given by the sum of the connections induced on V and $V^\perp$ by $\hat{d}$.

Definition 2. *A non-degenerate rank* 4 *subbundle* V *of* $\underline{\mathbb{C}}^{n+2}$ *is said to be a* $\hat{d}$-central sphere congruence *of a* $\hat{d}$*-surface* $(\Lambda^{1,0},\Lambda^{0,1})$ *if* $\Lambda^{1,0}+\Lambda^{0,1}\subset V$ *and*

$$(\mathcal{N}_V^{\hat{d}})^{1,0}\Lambda^{0,1} = 0 = (\mathcal{N}_V^{\hat{d}})^{0,1}\Lambda^{1,0}.$$

Remark 1. *Let* $(\Lambda^{1,0},\Lambda^{0,1})$ *be a surface,* $\sigma\neq 0$ *be a section of* Λ *and* z *be a holomorphic chart of* Σ*. Note that* $\Lambda^{1,0}+\Lambda^{0,1}\subset V$ *establishes* $(\mathcal{N}_V)_{|\Lambda}=0$ *and then* $\mathcal{N}_V^{1,0}\Lambda^{0,1} = 0 = \mathcal{N}_V^{0,1}\Lambda^{1,0}$ *reads* $\sigma_{z\bar{z}}\in\Gamma V$*. Hence, generically (if* $\sigma\wedge\sigma_z\neq 0\neq\sigma\wedge\sigma_{\bar{z}}$*),* $\Lambda^{1,0}$, $\Lambda^{0,1}$ *and* V *are all determined by* Λ*:* $\Lambda^{1,0}=\langle\sigma,\sigma_z\rangle$, $\Lambda^{0,1}=\langle\sigma,\sigma_{\bar{z}}\rangle$ *and* $V=\langle\sigma,\sigma_z,\sigma_{\bar{z}},\sigma_{z\bar{z}}\rangle$*. In particular, the complexification of the central sphere congruence of a real surface* Λ *is the unique central sphere congruence of the corresponding surface* $(\Lambda^{1,0},\Lambda^{0,1})$*.*

For further reference:

Lemma 6. *Suppose* V *is a* $\hat{d}$*-central sphere congruence of a* $\hat{d}$*-surface* $(\Lambda^{1,0},\Lambda^{0,1})$*. Then*

$$(\mathcal{D}_V^{\hat{d}})^{1,0}\Gamma\Lambda^{1,0}\subset\Omega^{1,0}\Lambda^{1,0}, \quad (\mathcal{D}_V^{\hat{d}})^{0,1}\Gamma\Lambda^{0,1}\subset\Omega^{0,1}\Lambda^{0,1}.$$

Proof. First of all, observe that, as $\operatorname{rank}\Lambda^{1,0} = \frac{1}{2}\operatorname{rank} V = \operatorname{rank}\Lambda^{0,1}$, the isotropy of both $\Lambda^{1,0}$ and $\Lambda^{0,1}$ establishes their maximal isotropy in V.

Write $\Lambda^{1,0} = \langle \sigma, \tau \rangle$, with $\sigma \in \Gamma\Lambda$. By definition of $\hat{d}$-surface and of $\hat{d}$-central sphere congruence, $(\mathcal{D}_V^{\hat{d}})^{1,0}\sigma = \pi_V \circ \hat{d}^{1,0} \circ \pi_V \sigma \in \Omega^{1,0}\Lambda^{1,0}$. By the isotropy of $\Lambda^{1,0}$ and the fact that $\mathcal{D}_V^{\hat{d}}$ is a metric connection, it follows that $((\mathcal{D}_V^{\hat{d}})^{1,0}\tau, \sigma) = -(\tau, (\mathcal{D}_V^{\hat{d}})^{1,0}\sigma) = 0$, while $((\mathcal{D}_V^{\hat{d}})^{1,0}\tau, \tau) = \frac{1}{2}\, d^{1,0}(\tau,\tau) = 0$. We conclude that $(\mathcal{D}_V^{\hat{d}})^{1,0}\tau \perp \Lambda^{1,0}$ and, therefore, that it takes values in $\Lambda^{1,0}$. A similar argument establishes $(\mathcal{D}_V^{\hat{d}})^{0,1}\Gamma\Lambda^{0,1} \subset \Omega^{0,1}\Lambda^{0,1}$. □

Definition 3. *A non-degenerate bundle $V \subset \underline{\mathbb{C}}^{n+2}$ is said to be* $\hat{d}$-constrained harmonic *if there exists a 1-form q with values in $\wedge^2 V \oplus \wedge^2 V^\perp$ such that, for each $\lambda \in \mathbb{C}\backslash\{0\}$, the metric connection*

$$\hat{d}_V^{\lambda,q} := \mathcal{D}_V^{\hat{d}} + \lambda^{-1}(\mathcal{N}_V^{\hat{d}})^{1,0} + \lambda(\mathcal{N}_V^{\hat{d}})^{0,1} + (\lambda^{-2}-1)q^{1,0} + (\lambda^2-1)q^{0,1},$$

on $\underline{\mathbb{C}}^{n+2}$, is flat. In this case, we say that V is $(q,\hat{d})$-constrained harmonic *or, in the case $q=0$,* $\hat{d}$-harmonic. *In the particular case of $\hat{d} = d$, V a real bundle and q a real form, we say that V is a* real q-constrained harmonic bundle.

Definition 4. *A $\hat{d}$-surface $(\Lambda^{1,0}, \Lambda^{0,1})$ is said to be* $\hat{d}$-constrained Willmore *if it admits a $(q,\hat{d})$-constrained harmonic $\hat{d}$-central sphere congruence with*

$$q^{1,0} \in \Omega^{1,0}(\wedge^2\Lambda^{0,1}), \quad q^{0,1} \in \Omega^{0,1}(\wedge^2\Lambda^{1,0}). \tag{13}$$

If $q = 0$, we say that $(\Lambda^{1,0}, \Lambda^{0,1})$ is $\hat{d}$-Willmore. *In the particular case of $\hat{d} = d$, $(\Lambda^{1,0}, \Lambda^{0,1})$ a real surface and q a real form, we say that $(\Lambda^{1,0}, \Lambda^{0,1})$ is a a* real q-constrained Willmore surface.

From Theorem 5 and Proposition 4, it follows that the real constrained Willmore surface condition is preserved under the correspondence

$$(\Lambda^{1,0}, \Lambda^{0,1}) \longleftrightarrow \Lambda^{1,0} \cap \Lambda^{0,1}$$

for real surfaces $(\Lambda^{1,0}, \Lambda^{0,1})$, with preservation of multipliers:

Theorem 7. *Suppose $(\Lambda^{1,0}, \Lambda^{0,1})$ is a real surface. Then Λ is constrained Willmore if and only if S is q-constrained harmonic, for some real 1-form q with $q^{1,0} \in \Omega^{1,0}(\wedge^2\Lambda^{0,1})$.*

3. TRANSFORMATIONS OF CONSTRAINED HARMONIC BUNDLES AND CONSTRAINED WILLMORE SURFACES

Fix a conformal structure $\mathcal{C}$ on Σ and consider the corresponding complex structure on Σ. Let Ad denote the adjoint representation of the orthogonal group on the orthogonal algebra. Note that, given $T \in O(\mathbb{R}^{n+1,1})$ and $u, v \in \mathbb{R}^{n+1,1}$, $\mathrm{Ad}_T(u \wedge v) = Tu \wedge Tv$.

Let V be a non-degenerate subbundle of $\underline{\mathbb{C}}^{n+2}$ and π_V and $\pi_{V^\perp}$ denote the orthogonal projections of $\underline{\mathbb{C}}^{n+2}$ onto V and $V^\perp$, respectively, and ρ

denote reflection across V, $\rho = \pi_V - \pi_{V^\perp}$. Let $(\Lambda^{1,0}, \Lambda^{0,1})$ be a surface admitting V as a central sphere congruence. As usual, we write Λ for $\Lambda^{1,0} \cap \Lambda^{0,1}$. Suppose V is q-constrained harmonic for some $q \in \Omega^1(\wedge^2 V \oplus \wedge^2 V^\perp)$ satisfying conditions (13).

3.1. **Spectral deformation.** For each $\lambda \in \mathbb{C}\backslash\{0\}$, the flatness of the metric connection $d_V^{\lambda,q}$ on $\underline{\mathbb{C}}^{n+2}$ establishes the existence of an isometry

$$\phi_V^{\lambda,q} : (\underline{\mathbb{C}}^{n+2}, d_V^{\lambda,q}) \to (\underline{\mathbb{C}}^{n+2}, d)$$

of bundles, preserving connections, defined on a simply connected component of Σ and unique up to a Möbius transformation.

Lemma 8. *Let $\hat{d}$ be a flat metric connection on $\underline{\mathbb{C}}^{n+2}$ and $\phi : (\underline{\mathbb{C}}^{n+2}, \hat{d}) \to (\underline{\mathbb{C}}^{n+2}, d)$ be an isometry of bundles, preserving connections. Then V is $(q, \hat{d})$-constrained harmonic, for some q, if and only if ϕV is $\mathrm{Ad}_\phi q$-constrained harmonic.*

Proof. It is immediate from the fact that

$$\mathcal{D}_{\phi V} = \phi \circ \mathcal{D}_V^{\hat{d}} \circ \phi^{-1}, \ \ \mathcal{N}_{\phi V} = \phi \mathcal{N}_V^{\hat{d}} \phi^{-1} \tag{14}$$

and, therefore, $d_{\phi V}^{\lambda,q} = \phi \circ \hat{d}_V^{\lambda, \mathrm{Ad}_{\phi^{-1}} q} \circ \phi^{-1}$. □

Set

$$q_\lambda := \lambda^{-2} q^{1,0} + \lambda^2 q^{0,1},$$

for $\lambda \in \mathbb{C}\backslash\{0\}$. The fact that q takes values in $\wedge^2 V \oplus \wedge^2 V^\perp$ establishes, in particular, $\mathcal{D}_V^{d_V^{\lambda,q}} = \mathcal{D}_V + (\lambda^{-2} - 1) q^{1,0} + (\lambda^{0,1} - 1) q^{0,1}$, whereas $\mathcal{N}_V^{d_V^{\lambda,q}} = \lambda^{-1} \mathcal{N}_V^{1,0} + \lambda \mathcal{N}_V^{0,1}$, and, therefore, $(d_V^{\lambda,q})_V^{\mu, q_\lambda} = d_V^{\lambda\mu, q}$, for all $\lambda, \mu \in \mathbb{C}\backslash\{0\}$. From the flatness of $d_V^{\lambda,q}$, for all $\lambda \in \mathbb{C}\backslash\{0\}$, we conclude that of $(d_V^{\lambda,q})_V^{\mu,q_\lambda}$, for all $\lambda, \mu \in \mathbb{C}\backslash\{0\}$ and, therefore, that V is $d_V^{\lambda,q}$-constrained harmonic, for all $\lambda \in \backslash\mathbb{C}\{0\}$. We define then a spectral deformation of V into new constrained harmonic bundles by setting, for each λ in $\mathbb{C}\{0\}$,

$$V_q^\lambda := \phi_V^{\lambda,q} V.$$

Theorem 9. *V_q^λ is $\mathrm{Ad}_{\phi_q^\lambda} q_\lambda$-constrained harmonic, for each $\lambda \in \mathbb{C}\{0\}$.*

A deformation on the level of constrained Willmore surfaces follows:

Theorem 10. *For each $\lambda \in \mathbb{C}\backslash\{0\}$, $(\phi_V^{\lambda,q} \Lambda^{1,0}, \phi_V^{\lambda,q} \Lambda^{0,1})$ is a $\mathrm{Ad}_{\phi_q^\lambda}(q_\lambda)$-constrained Willmore surface, admitting V_q^λ as a central sphere congruence. Furthermore, if $(\Lambda^{1,0}, \Lambda^{0,1})$ is a real constrained Willmore surface, then so is*

$$\Lambda_q^\lambda := \phi_V^{\lambda,q} \Lambda,$$

for all $\lambda \in S^1$.

Proof. By (13), together with the isotropy of $\Lambda^{i,j}$, for $i \neq j \in \{0,1\}$, we have $q\Lambda = 0$. On the other hand, the centrality of V with respect to $(\Lambda^{1,0}, \Lambda^{0,1})$ gives $\mathcal{N}_V \Lambda = \pi_{V^\perp} \circ d\Lambda = 0$. Hence

$$(d_V^{\lambda,q})_{|\Gamma\Lambda} = d_{|\Gamma\Lambda} \tag{15}$$

and we conclude that $(\phi_V^{\lambda,q}\,\Lambda^{1,0}, \phi_V^{\lambda,q}\,\Lambda^{0,1})$ is still a surface. Suppose, furthermore, that $(\Lambda^{1,0}, \Lambda^{0,1})$ is a real q-constrained Willmore surface. Given $\lambda \in S^1$, d_q^λ is real, so that we can choose $\phi_V^{\lambda,q}$ to be real, in which case $\overline{\phi_V^{\lambda,q}\Lambda} = \phi_V^{\lambda,q}\Lambda$, Λ_q^λ is a real surface. It is obvious, on the other hand, that as V is a central sphere congruence of $(\Lambda^{1,0}, \Lambda^{0,1})$, $\phi_V^{\lambda,q}\,V$ is a central sphere congruence of $(\phi_V^{\lambda,q}\,\Lambda^{1,0}, \phi_V^{\lambda,q}\,\Lambda^{0,1})$. Theorem 9 completes the proof. □

Note that spectral deformation corresponding to the zero multiplier preserves the class of Willmore surfaces. This spectral deformation of real (constrained) Willmore surfaces coincides, up to reparametrization, with the one presented in [8], in terms of the *Schwarzian derivative* and the *Hopf differential*.

An alternative perspective on this spectral deformation of (constrained) harmonic bundles and Willmore surfaces is that of a change of flat connection on $\underline{\mathbb{C}}^{n+2}$: if $V \subset (\underline{\mathbb{C}}^{n+2}, d)$ is constrained harmonic, then so is $V \subset (\underline{\mathbb{C}}^{n+2}, d_V^{\lambda,q})$, as well as, if $(\Lambda^{1,0}, \Lambda^{0,1})$ is constrained Willmore [with respect to d] then it is still constrained Willmore with respect to $d_V^{\lambda,q}$, for all $\lambda \in \mathbb{C}\backslash\{0\}$. In the real case, this is the interpretation of loop group theory in [5].

3.2. **Dressing action.** We use a version of the dressing action theory of Terng and Uhlenbeck [19] to build transforms of V into new constrained harmonic bundles and thereafter transforms of $(\Lambda^{1,0}, \Lambda^{0,1})$ into new constrained Willmore surfaces. For that, we give conditions on a dressing $r(\lambda) \in \Gamma(O(\underline{\mathbb{C}}^{n+2}))$ such that the gauging $r(\lambda) \circ d_V^{\lambda,q} \circ r(\lambda)^{-1}$ of $d_V^{\lambda,q}$ by $r(\lambda)$ establishes the constrained harmonicity of some bundle $\hat{V}$ from the constrained harmonicity of V.

The $\mathcal{D}_V$-parallelness of V and $V^\perp$, together with the fact that $\mathcal{N}_V$ intertwines V and $V^\perp$, whereas q preserves them, makes clear that

$$d_V^{-\lambda,q} = \rho \circ d_V^{\lambda,q} \circ \rho^{-1}, \tag{16}$$

for $\lambda \in \mathbb{C}\backslash\{0\}$. Suppose we have $r(\lambda) \in \Gamma(O(\underline{\mathbb{C}}^{n+2}))$ such that $\lambda \mapsto r(\lambda)$ is rational in λ, r is holomorphic and invertible at $\lambda = 0$ and $\lambda = \infty$ and twisted in the sense that

$$\rho\, r(\lambda)\, \rho^{-1} = r(-\lambda), \tag{17}$$

for $\lambda \in \mathrm{dom}(r)$. In particular, it follows that both $r(0)$ and $r(\infty)$ commute with ρ, and, therefore, that

$$r(0)_{|_V}, r(\infty)_{|_V} \in \Gamma(O(V)). \tag{18}$$

Define $\hat{q} \in \Omega^1(\wedge^2 V \oplus \wedge^2 V^\perp)$ by setting

$$\hat{q}^{1,0} := \mathrm{Ad}_{r(0)} q^{1,0}, \quad \hat{q}^{\,0,1} := \mathrm{Ad}_{r(\infty)} q^{0,1}.$$

Define a new family of metric connections on $\underline{\mathbb{C}}^{n+2}$ by setting

$$\hat{d}_V^{\lambda,\hat{q}} := r(\lambda) \circ d_V^{\lambda,q} \circ r(\lambda)^{-1}.$$

Suppose that there exists a holomorphic extension of $\lambda \mapsto \hat{d}_V^{\lambda,\hat{q}}$ to $\lambda \in \mathbb{C}\backslash\{0\}$ through metric connections on $\underline{\mathbb{C}}^{n+2}$. We shall see later how to construct such $r = r(\lambda)$, but assume, for the moment, that we have got one. In that case, as we, crucially, verify next, the notation $\hat{d}_V^{\lambda,\hat{q}}$ is not merely formal:

Proposition 11.

$$\hat{d}_V^{\lambda,\hat{q}} = \mathcal{D}_V^{\hat{d}} + \lambda^{-1}(\mathcal{N}_V^{\hat{d}})^{1,0} + \lambda(\mathcal{N}_V^{\hat{d}})^{0,1} + (\lambda^{-2} - 1)\hat{q}^{1,0} + (\lambda^2 - 1)\hat{q}^{0,1},$$

for the flat metric connection $\hat{d} := \hat{d}_V^{1,\hat{q}} = \lim_{\lambda\to 1} r(\lambda) \circ d_V^{\lambda,q} \circ r(\lambda)^{-1}$ *and* $\lambda \in \mathbb{C}\backslash\{0\}$.

Proof. The fact that r is holomorphic and invertible at $\lambda = 0$ and $(d_V^{\lambda,q})^{0,1} = \mathcal{D}_V^{0,1} + \lambda\mathcal{N}_V^{0,1} + (\lambda^2 - 1)q^{0,1}$ is holomorphic on $\mathbb{C}$ establishes that the connection $(\hat{d}_V^{\lambda,\hat{q}})^{0,1} = r(\lambda) \circ (d_V^{\lambda,q})^{0,1} \circ r(\lambda)^{-1}$, which admits a holomorphic extension to $\lambda \in \mathbb{C}\backslash\{0\}$, admits, furthermore, a holomorphic extension to $\lambda \in \mathbb{C}$. Thus, locally, $(\hat{d}_V^{\lambda,\hat{q}})^{0,1} = A_0^{0,1} + \sum_{i\geq 1} \lambda^i A_i^{0,1}$, with A_0 connection and $A_i \in \Omega^1(o(\underline{\mathbb{C}}^{n+2}))$, for all i. Considering then limits of

$$\lambda^{-2}A_0^{0,1} + \sum_{i\geq 1} \lambda^{i-2} A_i^{0,1} = r(\lambda) \circ (\lambda^{-2}\mathcal{D}_V^{0,1} + \lambda^{-1}\mathcal{N}_V^{0,1} + (1 - \lambda^{-2})q^{0,1}) \circ r(\lambda)^{-1},$$

when λ goes to infinity, we get $A_2^{0,1} + \lim_{\lambda\to\infty} \sum_{i\geq 3} \lambda^{i-2} A_i^{0,1} = \mathrm{Ad}_{r(\infty)}\, q^{0,1}$, which shows that $A_i^{0,1} = 0$, for all $i \geq 3$, and that $A_2^{0,1} = \hat{q}^{0,1}$. Considering now limits of

$$A_0^{0,1} + \lambda A_1^{0,1} + \lambda^2 \hat{q}^{0,1} = r(\lambda) \circ (\mathcal{D}_V^{0,1} + \lambda\mathcal{N}_V^{0,1} + (\lambda^2 - 1)q^{0,1}) \circ r(\lambda)^{-1},$$

when λ goes to 0, we conclude that $A_0^{0,1} = r(0) \circ (\mathcal{D}_V^{0,1} - q^{0,1}) \circ r(0)^{-1}$ and, therefore, that $(\hat{d}_V^{\lambda,\hat{q}})^{0,1} = r(0) \circ (\mathcal{D}_V^{0,1} - q^{0,1}) \circ r(0)^{-1} + \lambda A_1^{0,1} + \lambda^2 \hat{q}^{0,1}$. As for $(\hat{d}_V^{\lambda,\hat{q}})^{1,0} = r(\lambda) \circ (\mathcal{D}_V^{1,0} + \lambda^{-1}\mathcal{N}^{1,0} + (\lambda^{-2} - 1)q^{1,0}) \circ r(\lambda)^{-1}$, which has a pole at $\lambda = 0$, we have, for λ away from 0,

$$\sum_{i\geq 1} \lambda^{-i} A_{-i}^{1,0} + A_0^{1,0} + \sum_{i\geq 1} \lambda^i A_i^{1,0} = r(\lambda) \circ (\mathcal{D}_V^{1,0} + \lambda^{-1}\mathcal{N}^{1,0} + (\lambda^{-2} - 1)q^{1,0}) \circ r(\lambda)^{-1}, \tag{19}$$

with $A_{-i}^{1,0} \in \Omega^1(o(\underline{\mathbb{C}}^{n+2}))$, for all $i \geq 1$. Considering limits of (19) when λ goes to infinity, shows that $A_i^{1,0} = 0$, for all $i \geq 1$, and that $A_0^{1,0} =$

$r(\infty)\circ(\mathcal{D}_V^{1,0}-q^{1,0})\circ r(\infty)^{-1}$. Multiplying then both members of equation (19) by λ^2 and considering limits when λ goes to 0, we conclude that $A_{-2}^{1,0}=\hat{q}^{1,0}$ and that $A_{-i}^{1,0}=0$, for all $i\geq 3$, and, ultimately, that $(\hat{d}_V^{\lambda,\hat{q}})^{1,0}=r(\infty)\circ(\mathcal{D}_V^{1,0}-q^{1,0})\circ r(\infty)^{-1}+\lambda^{-1}A_{-1}^{1,0}+\lambda^{-2}\hat{q}^{1,0}$. Thus

$$\begin{aligned}\hat{d}_V^{\lambda,\hat{q}} \quad = \quad & r(0)\circ(\mathcal{D}_V^{0,1}-q^{0,1}+q^{1,0})\circ r(0)^{-1}\\ & +r(\infty)\circ(\mathcal{D}_V^{1,0}-q^{1,0}+q^{0,1})\circ r(\infty)^{-1}\\ & +\lambda^{-1}A_{-1}^{1,0}+\lambda A_1^{0,1}+(\lambda^{-2}-1)\hat{q}^{1,0}+(\lambda^2-1)\hat{q}^{0,1},\end{aligned}$$

for $\lambda\in\mathbb{C}\backslash\{0\}$, and, in particular,

$$\hat{d}=r(0)\circ(\mathcal{D}_V^{0,1}-q^{0,1}+q^{1,0})\circ r(0)^{-1}+r(\infty)\circ(\mathcal{D}_V^{1,0}-q^{1,0}+q^{0,1})\circ r(\infty)^{-1}+A_{-1}^{1,0}+A_1^{0,1}.$$

The fact that $r(0)$ and $r(\infty)$ (and so $r(0)^{-1}$ and $r(\infty)^{-1}$), as well as q, preserve V and $V^\perp$, together with the $\mathcal{D}_V$-parallelness of V and of $V^\perp$, shows that $\hat{d}-(A_{-1}^{1,0}+A_1^{0,1})$ preserves $\Gamma(V)$ and $\Gamma(V^\perp)$. On the other hand, equations (16) and (17) combine to give $\hat{d}_V^{-\lambda,\hat{q}}=\rho\circ\hat{d}_V^{\lambda,\hat{q}}\circ\rho^{-1}$, for all $\lambda\in\mathbb{C}\backslash\{0\}$ away from the poles of r and then, by continuity, on all of $\mathbb{C}\backslash\{0\}$. The particular case of $\lambda=1$ gives $\rho(A_{-1}^{1,0}+A_1^{0,1})_{|V}=-(A_{-1}^{1,0}+A_1^{0,1})_{|V}$ and $\rho(A_{-1}^{1,0}+A_1^{0,1})_{|V^\perp}=-(A_{-1}^{1,0}+A_1^{0,1})_{|V^\perp}$, showing that $A_{-1}^{1,0}+A_1^{0,1}\in\Omega^1(V\wedge V^\perp)$. We conclude that

$$r(0)\circ(\mathcal{D}_V^{0,1}-q^{0,1}+q^{1,0})\circ r(0)^{-1}+r(\infty)\circ(\mathcal{D}_V^{1,0}-q^{1,0}+q^{0,1})\circ r(\infty)^{-1}=\mathcal{D}_V^{\hat{d}} \tag{20}$$

and $A_{-1}^{1,0}=(\mathcal{N}_V^{\hat{d}})^{1,0}$, $A_1^{0,1}=(\mathcal{N}_V^{\hat{d}})^{0,1}$, completing the proof. □

The flatness of $d_V^{\lambda,q}$ for all $\lambda\in\mathbb{C}\backslash\{0\}$ establishes that of $\hat{d}_V^{\lambda,\hat{q}}$, for all non-zero λ away from the poles of r and then, by continuity, for all $\lambda\in\mathbb{C}\backslash\{0\}$. By Proposition 11, we conclude that V is $(\hat{q},\hat{d})$-constrained harmonic. Suppose $1\in\mathrm{dom}(r)$. By Lemma 8, it follows that:

Theorem 12. *$r(1)^{-1}V$ is a $\mathrm{Ad}_{r(1)^{-1}}\,\hat{q}$-constrained harmonic bundle.*

Note that this transformation preserves the harmonicity condition.

A transformation on the level of constrained harmonic surfaces follows, with some extra condition, as follows. Set

$$\hat{\Lambda}^{1,0}:=r(\infty)\Lambda^{1,0},\quad \hat{\Lambda}^{0,1}:=r(0)\Lambda^{0,1}$$

and

$$\hat{\Lambda}=\hat{\Lambda}^{1,0}\cap\hat{\Lambda}^{0,1}.$$

Suppose, furthermore, that

$$\det r(0)_{|V}=\det r(\infty)_{|V}. \tag{21}$$

Then:

Theorem 13. *$(r(1)^{-1}\hat{\Lambda}^{1,0},r(1)^{-1}\hat{\Lambda}^{0,1})$ is a $\mathrm{Ad}_{r(1)^{-1}}\hat{q}$-constrained Willmore surface admitting $r(1)^{-1}V$ as a central sphere congruence.*

Proof. First of all, note that, by (13), $\hat{q}^{i,j} \in \Omega^{i,j}(\wedge^2 \hat{\Lambda}^{j,i})$ and, therefore, $(\mathrm{Ad}_{r(1)^{-1}}\hat{q})^{i,j} \in \Omega^{i,j}(\wedge^2 r(1)^{-1}\hat{\Lambda}^{j,i})$, for $i \neq j \in \{0,1\}$. In the light of Theorem 12, we are left to verify that $(r(1)^{-1}\hat{\Lambda}^{1,0}, r(1)^{-1}\hat{\Lambda}^{0,1})$ is a surface admitting $r(1)^{-1}V$ as a central sphere congruence.

The fact that $\Lambda^{1,0}$ and $\Lambda^{0,1}$ are rank 2 isotropic subbundles of V ensures that so are $\Lambda^{\hat{1},0}$ and $\Lambda^{\hat{0},1}$, as $r(0)$ and $r(\infty)$ are orthogonal transformations and preserve V. To see that $\hat{\Lambda}$ is rank 1, we use some well-known facts about the Grassmannian $\mathcal{G}_W$ of isotropic 2-planes in a complex 4-dimensional space W: it has two components, each an orbit of the special orthogonal group $SO(W)$, intertwined by the action of elements of $O(W)\backslash SO(W)$, and for which any element intersects any element of the other component in a line while distinct elements of the same component have trivial intersection. Since $\operatorname{rank}\Lambda = 1$, $\Lambda^{1,0}_p$ and $\Lambda^{0,1}_p$ lie in different components of $\mathcal{G}_{V_p}$ and the hypothesis (21) ensures that the same is true of $\hat{\Lambda}^{1,0}_p$ and $\hat{\Lambda}^{0,1}_p$, for all p.

We are left to verify that

$$\hat{d}^{1,0}\Gamma(\hat{\Lambda}) \subset \Omega^1(\hat{\Lambda}^{1,0}), \quad \hat{d}^{0,1}\Gamma(\hat{\Lambda}) \subset \Omega^1(\hat{\Lambda}^{0,1}) \tag{22}$$

and that (recall equation (14))

$$(\mathcal{N}^{\hat{d}}_V)^{1,0}\hat{\Lambda}^{0,1} = 0 = (\mathcal{N}^{\hat{d}}_V)^{0,1}\hat{\Lambda}^{1,0}. \tag{23}$$

Equation (23) forces $\mathcal{N}^{\hat{d}}_V\hat{\Lambda} = 0$, in which situation, (22) reads

$$(\mathcal{D}^{\hat{d}}_V)^{1,0}\Gamma(\hat{\Lambda}) \subset \Omega^1(\hat{\Lambda}^{1,0}), \quad (\mathcal{D}^{\hat{d}}_V)^{0,1}\Gamma(\hat{\Lambda}) \subset \Omega^1(\hat{\Lambda}^{0,1}),$$

which, in its turn, follows from

$$(\mathcal{D}^{\hat{d}}_V)^{1,0}\Gamma(\hat{\Lambda}^{1,0}) \subset \Omega^1(\hat{\Lambda}^{1,0}), \quad (\mathcal{D}^{\hat{d}}_V)^{0,1}\Gamma(\hat{\Lambda}^{0,1}) \subset \Omega^1(\hat{\Lambda}^{0,1}). \tag{24}$$

It is (23) and (24) that we shall establish.

First of all, note that, according to (20),

$$(\mathcal{D}^{\hat{d}}_V)^{1,0} = r(\infty) \circ (\mathcal{D}^{1,0}_V - q^{1,0}) \circ r(\infty)^{-1} + \hat{q}^{1,0}$$

and

$$(\mathcal{D}^{\hat{d}}_V)^{0,1} = r(0) \circ (\mathcal{D}^{0,1}_V - q^{0,1}) \circ r(0)^{-1} + \hat{q}^{0,1}.$$

Now $q^{1,0}$ takes values in $\Lambda \wedge \Lambda^{0,1}$, so $q^{1,0}\Lambda^{1,0} \subset \Lambda \subset \Lambda^{1,0}$, by the isotropy of $\Lambda^{1,0}$. On the other hand, since $\operatorname{rank}\hat{\Lambda} = 1$, we have $\wedge^2\hat{\Lambda}^{0,1} = \hat{\Lambda} \wedge \hat{\Lambda}^{0,1}$ and, therefore, $\hat{q}^{1,0}\hat{\Lambda}^{1,0} \subset \hat{\Lambda} \subset \hat{\Lambda}^{1,0}$. Together with Lemma 6, this establishes the $(1,0)$-part of (24). A similar argument establishes the $(0,1)$-part of it.

Finally, we establish (23). According to Proposition 11,

$$\begin{aligned}
(\mathcal{N}^{\hat{d}}_V)^{1,0} &= \lim\nolimits_{\lambda\to 0} \lambda((\hat{d}^{\lambda,\hat{q}}_V)^{1,0} - (\mathcal{D}^{\hat{d}}_V)^{1,0} - (\lambda^{-2} - 1)\hat{q}^{1,0}) \\
&= \lim\nolimits_{\lambda\to 0} \lambda((\hat{d}^{\lambda,\hat{q}}_V)^{1,0} - \lambda^{-2}\hat{q}^{1,0}) \\
&= \lim\nolimits_{\lambda\to 0} (r(\lambda) \circ (\lambda\, (d^{\lambda,q}_V)^{1,0}) \circ r(\lambda)^{-1} - \lambda^{-1}\mathrm{Ad}_{r(0)}q^{1,0}) \\
&= \mathrm{Ad}_{r(0)}\mathcal{N}^{1,0}_V + \lim\nolimits_{\lambda\to 0} \frac{1}{\lambda}(\mathrm{Ad}_{r(\lambda)} - \mathrm{Ad}_{r(0)})q^{1,0}.
\end{aligned}$$

so that

$$(\mathcal{N}_V^{\hat d})^{1,0} = \mathrm{Ad}_{r(0)}\mathcal{N}_V^{1,0} + \frac{d}{d\lambda}_{|\lambda=0}\mathrm{Ad}_{r(\lambda)}q^{1,0}; \tag{25}$$

and, similarly,

$$\begin{aligned}(\mathcal{N}_V^{\hat d})^{0,1} &= \lim\nolimits_{\lambda\to\infty}\lambda^{-1}((\hat d_V^{\lambda,\hat q})^{0,1} - (\mathcal{D}_V^{\hat d})^{0,1} - (\lambda^2-1)\hat q^{0,1})\\ &= \mathrm{Ad}_{r(\infty)}\mathcal{N}_V^{0,1} + \lim\nolimits_{\lambda\to\infty}(r(\lambda)\circ\lambda q^{0,1}\circ r(\lambda)^{-1} - \lambda\mathrm{Ad}_{r(\infty)}q^{0,1})\end{aligned}$$

and, therefore,

$$(\mathcal{N}_V^{\hat d})^{0,1} = \mathrm{Ad}_{r(\infty)}\mathcal{N}_V^{0,1} + \frac{d}{d\lambda}_{|\lambda=0}\mathrm{Ad}_{r(\lambda^{-1})}q^{0,1}. \tag{26}$$

Furthermore, by (25),

$$(\mathcal{N}_V^{\hat d})^{1,0} = \mathrm{Ad}_{r(0)}(\mathcal{N}^{1,0} + [r(0)^{-1}\frac{d}{d\lambda}_{|\lambda=0}r(\lambda), q^{1,0}]).$$

The centrality of V with respect to $(\Lambda^{1,0},\Lambda^{0,1})$ establishes, in particular, $\mathcal{N}_V^{1,0}\Lambda^{0,1} = 0$, whilst the isotropy of $\Lambda^{0,1}$ ensures, in particular, that $q^{1,0}\Lambda^{0,1} = 0$. Hence

$$\mathrm{Ad}_{r(0)}(\mathcal{N}^{1,0} + r(0)^{-1}\frac{d}{d\lambda}_{|\lambda=0}r(\lambda)\,q^{1,0})\hat\Lambda^{0,1} = 0.$$

On the other hand, differentiation of $r(\lambda)^{-1} = \rho\, r(-\lambda)^{-1}\rho$, derived from equation (17), gives

$$-r(\lambda)^{-1}\frac{d}{dk}_{|k=\lambda}r(k)\,r(\lambda)^{-1} = \rho\, r(-\lambda)^{-1}\frac{d}{dk}_{|k=-\lambda}r(k)\,r(-\lambda)^{-1}\rho,$$

or, equivalently,

$$\rho\, r(\lambda)^{-1}\frac{d}{dk}_{|k=\lambda}r(k)\rho = -r(-\lambda)^{-1}\frac{d}{dk}_{|k=-\lambda}r(k)\,r(-\lambda)^{-1}\rho\, r(\lambda)\rho,$$

and, therefore, yet again by equation (17),

$$\rho\, r(\lambda)^{-1}\frac{d}{dk}_{|k=\lambda}r(k)\rho = -r(-\lambda)^{-1}\frac{d}{dk}_{|k=-\lambda}r(k).$$

Evaluation at $\lambda = 0$ shows then that

$$\rho\, r(0)^{-1}\frac{d}{d\lambda}_{|\lambda=0}r(\lambda)\rho = -r(0)^{-1}\frac{d}{d\lambda}_{|\lambda=0}r(\lambda).$$

Equivalently,

$$r(0)^{-1}\frac{d}{d\lambda}_{|\lambda=0}r(\lambda) \in \Gamma(V\wedge V^{\perp}). \tag{27}$$

Since $qV^{\perp} = 0$, we conclude that $Ad_{r(0)}(q^{1,0}r(0)^{-1}\frac{d}{d\lambda}_{|\lambda=0}r(\lambda))\hat\Lambda^{0,1} = 0$ and, ultimately, that $(\mathcal{N}_V^{\hat d})^{1,0}\hat\Lambda^{0,1} = 0$. A similar argument near $\lambda = \infty$ establishes $(\mathcal{N}_V^{\hat d})^{0,1}\hat\Lambda^{1,0} = 0$, completing the proof. □

3.3. **Bäcklund transformation.** We now construct $r = r(\lambda)$ satisfying the hypothesis of the previous section. As the philosophy underlying the work of C.-L. Terng and K. Uhlenbeck [21] suggests, we consider linear fractional transformations. As we shall see, a two-step process will produce a desired r.

Given $\alpha \in \mathbb{C}\backslash\{-1,0,1\}$ and L a null line subbundle of $\underline{\mathbb{C}}^{n+2}$ such that, locally, $\rho L \cap L^{\perp} = \{0\}$, set

$$p_{\alpha,L}(\lambda) := I \begin{cases} \frac{\alpha-\lambda}{\alpha+\lambda} & \text{on}\, L \\ 1 & \text{on}\, (L\oplus\rho L)^{\perp} \\ \frac{\alpha+\lambda}{\alpha-\lambda} & \text{on}\, \rho L \end{cases}$$

and

$$q_{\alpha,L}(\lambda) := I \begin{cases} \frac{\lambda-\alpha}{\lambda+\alpha} & \text{on}\, L \\ 1 & \text{on}\, (L\oplus\rho L)^{\perp} \\ \frac{\lambda+\alpha}{\lambda-\alpha} & \text{on}\, \rho L \end{cases} ,$$

for $\lambda \in \mathbb{C}\backslash\{\pm\alpha\}$, defining in this way maps $p_{\alpha,L}, q_{\alpha,L} : \mathbb{C}\backslash\{\pm\alpha\} \to \Gamma(O(\underline{\mathbb{C}}^{n+2}))$ that, clearly, extend holomorphically to $\mathbb{P}^1\backslash\{\pm\alpha\}$, by setting

$$p_{\alpha,L}(\infty) :== I \begin{cases} -1 & \text{on}\, L \\ 1 & \text{on}\, (L\oplus\rho L)^{\perp} \\ -1 & \text{on}\, \rho L \end{cases}$$

and $q_{\alpha,L}(\infty) := I$. Obviously, $p_{\alpha,L}(\infty)$ and $q_{\alpha,L}(\infty)$ do not depend on α. For further reference, note that, for all $\lambda \in \mathbb{C}\backslash\{\pm\alpha, 0\}$, we have

$$p_{\alpha,L}(\lambda) = q_{\alpha^{-1},L}(\lambda^{-1}), \tag{28}$$

whilst $p_{\alpha,L}(0) = q_{\alpha,L}(\infty)$ and $p_{\alpha,L}(\infty) = q_{\alpha,L}(0)$.

The isometry $\rho = \rho^{-1}$ intertwines L and ρL and, therefore, preserves $(L \oplus \rho L)^{\perp}$, which makes clear that $\rho \circ p_{\alpha,L}(\lambda)$ and $p_{\alpha,L}(\lambda)^{-1} \circ \rho$ coincide in L, ρL and $(L\oplus\rho L)^{\perp}$ and, therefore,

$$\rho\, p_{\alpha,L}(\lambda)\rho = p_{\alpha,L}(\lambda)^{-1} = p_{\alpha,L}(-\lambda), \tag{29}$$

and, similarly,

$$\rho\, q_{\alpha,L}(\lambda)\rho = q_{\alpha,L}(\lambda)^{-1} = q_{\alpha,L}(-\lambda), \tag{30}$$

for all λ - both $p_{\alpha,L}$ and $q_{\alpha,L}$ are twisted in the sense of Section 3.2.

Since ρL is not orthogonal to L, $\rho L \neq L$, so L is not a subbundle of V and, therefore, $\operatorname{rank} V \cap (L \oplus \rho L) = 1$. We conclude that

$$p_{\alpha,L}(\infty)_{|V} = q_{\alpha,L}(0)_{|V} = I \begin{cases} -1 & \text{on}\, V\cap(L\oplus\rho L) \\ 1 & \text{on}\, V\cap(L\oplus\rho L)^{\perp} \end{cases}$$

has determinant $-1 \neq 1 = \det p_{\alpha,L}(0)_{|V} = \det q_{\alpha,L}(\infty)_{|V}$, so we cannot take $p_{\alpha,L} = r$ or $q_{\alpha,L} = r$ in the analysis of Section 3.2. However, we will be able to take $r = q_{\beta,\hat{L}}\, p_{\alpha,L}$, for suitable β and $\hat{L}$, as we shall see.

Now choose[1] $\alpha \in \mathbb{C}\backslash\{-1,0,1\}$ and L^α a $d_V^{\alpha,q}$-parallel null line subbundle of $\underline{\mathbb{C}}^{n+2}$ such that, locally,

$$\rho L^\alpha \cap (L^\alpha)^\perp = \{0\}. \tag{31}$$

Lemma 14. *There exists a holomorphic extension of*

$$\lambda \mapsto d^{\lambda,q}_{p_{\alpha,L^\alpha}} := p_{\alpha,L^\alpha}(\lambda) \circ d_V^{\lambda,q} \circ p_{\alpha,L^\alpha}(\lambda)^{-1}$$

to $\lambda \in \mathbb{C}\backslash\{0\}$ *through metric connections on* $\underline{\mathbb{C}}^{n+2}$*.*

Proof. We prove holomorphicity at $\lambda = \alpha$. Write $d_V^{\lambda,q} = d_V^{\alpha,q} + (\lambda-\alpha)A(\lambda)$, for $\lambda \in \mathbb{C}\backslash\{0,\alpha\}$, with $\lambda \mapsto A(\lambda) \in \Omega^1(o(\underline{\mathbb{C}}^{n+2}))$ holomorphic. Decompose $d_V^{\alpha,q} = D + \beta$ according to the decomposition

$$\underline{\mathbb{C}}^{n+2} = (L^\alpha \oplus \rho L^\alpha) \oplus (L^\alpha \oplus \rho L^\alpha)^\perp.$$

The fact that $d_V^{\alpha,q}$ is a metric connection establishes $d_V^{\alpha,q}\,\Gamma(\rho L^\alpha) \subset \Omega^1(\rho L^\alpha)^\perp$, as well as, in view of the $d_V^{\alpha,q}$-parallelness of L^α, $d_V^{\alpha,q}\,\Gamma((L^\alpha \oplus \rho L^\alpha)^\perp) \subset \Omega^1(L^\alpha)^\perp$. By (31), we conclude that the 1-form $\beta \in \Omega^1((L^\alpha \oplus \rho L^\alpha) \wedge (L^\alpha \oplus \rho L^\alpha)^\perp)$ takes values in $L^\alpha \wedge (L^\alpha \oplus \rho L^\alpha)^\perp$. Hence $p_{\alpha,L^\alpha}(\lambda) \circ \beta \circ p_{\alpha,L^\alpha}(\lambda)^{-1} = \frac{\alpha-\lambda}{\alpha+\lambda}\,\beta$. On the other hand, $p_{\alpha,L^\alpha}(\lambda) \circ D \circ p_{\alpha,L^\alpha}(\lambda)^{-1} = D$, as L^α, ρL^α and $(L^\alpha \oplus \rho L^\alpha)^\perp$ are all D-parallel. Thus

$$d^{\lambda,q}_{p_{\alpha,L}} = D + \frac{\alpha-\lambda}{\alpha+\lambda}\,\beta + (\lambda-\alpha)\,p_{\alpha,L^\alpha}(\lambda)\,A(\lambda)\,p_{\alpha,L^\alpha}(\lambda)^{-1}.$$

Lastly, note that, by the skew-symmetry of $A(\lambda)$, we have $A(\lambda)L^\alpha \subset (L^\alpha)^\perp$ and $A(\lambda)\rho L^\alpha \subset (\rho L^\alpha)^\perp$ and, therefore, by (31), $A(\lambda)L^\alpha \subset L^\alpha \oplus (L^\alpha \oplus \rho L^\alpha)^\perp$ and $A(\lambda)\rho L^\alpha \subset \rho L^\alpha \oplus (L^\alpha \oplus \rho L^\alpha)^\perp$. We conclude that $(\lambda-\alpha)\,\mathrm{Ad}_{p_{\alpha,L^\alpha}(\lambda)}\,A(\lambda)$ has at most a simple pole at $\lambda = -\alpha$ and, therefore, that $d^{\lambda,q}_{p_{\alpha,L}}$ is holomorphic at $\lambda = \alpha$. Furthermore, the fact that D is a metric connection establishes that so is $d^{\lambda,q}_{p_{\alpha,L}}$, in view of the skew-symmetry of $A(\lambda)$ and of β.

Holomorphicity at $\lambda = -\alpha$ can either be proved in the same way, having in consideration that the $d_V^{\alpha,q}$-parallelness of L^α establishes the $d_V^{-\alpha,q}$-parallelness of ρL^α, or by exploiting the symmetry $\lambda \mapsto -\lambda$. $\square$

Remark 2. 1) *The same argument establishes the existence of a holomorphic extension of*

$$\lambda \mapsto d^{\lambda,q}_{q_{\alpha,L^\alpha}} := q_{\alpha,L^\alpha}(\lambda) \circ d_V^{\lambda,q} \circ q_{\alpha,L^\alpha}(\lambda)^{-1}$$

to $\lambda \in \mathbb{C}\backslash\{0\}$ *through metric connections on* $\underline{\mathbb{C}}^{n+2}$*.*

2) *This argument uses nothing about the precise form of* $d_V^{\lambda,q}$*, only that it is holomorphic near* $\lambda = \pm\alpha$*.*

---

[1] Such a bundle L^α can be obtained by $d_V^{\alpha,q}$-parallel transport of $l_p^\alpha \in \mathbb{C}^{n+2}$, with l_p^α null and non-orthogonal to $\rho_p l_p^\alpha$, for some $p \in \Sigma$.

Now we can iterate the procedure starting with the connections $d^{\lambda,q}_{p_{\alpha,L^\alpha}}$: Choose $\beta \neq \pm\alpha$ in $\mathbb{C}\backslash\{-1,0,1\}$ and L^β a $d^{\beta,q}_V$-parallel null line subbundle of $\underline{\mathbb{C}}^{n+2}$. The fact that $p_{\alpha,L^\alpha} : (\underline{\mathbb{C}}^{n+2}, d^{\lambda,q}_V) \rightarrow (\underline{\mathbb{C}}^{n+2}, d^{\lambda,q}_{p_{\alpha,L^\alpha}})$ preserves connections establishes the $d^{\beta,q}_{p_{\alpha,L^\alpha}}$-parallelness of

$$\hat{L}^\beta_\alpha := p_{\alpha,L^\alpha}(\beta)L^\beta.$$

Choose L^β satisfying, furthermore, $\rho\hat{L}^\beta_\alpha \cap (\hat{L}^\beta_\alpha)^\perp = \{0\}$.[2] It follows that

$$\lambda \mapsto q_{\beta,\hat{L}^\beta_\alpha}(\lambda)\, p_{\alpha,L^\alpha}(\lambda) \circ d^{\lambda,q}_V \circ p_{\alpha,L^\alpha}(\lambda)^{-1}\, q_{\beta,\hat{L}^\beta_\alpha}(\lambda)^{-1}$$

admits a holomorphic extension to $\lambda \in \mathbb{C}\backslash\{0\}$ through metric connections on $\underline{\mathbb{C}}^{n+2}$ and, furthermore, that

$$r^* := q_{\beta,\hat{L}^\beta_\alpha} p_{\alpha,L^\alpha}$$

satisfies all the hypothesis of Section 3.2 on r. Set

$$q^* := \mathrm{Ad}_{r^*(1)^{-1}}(\mathrm{Ad}_{r^*(0)}q^{1,0} + \mathrm{Ad}_{r^*(\infty)}q^{0,1}).$$

Definition 5. *The q^*-constrained harmonic bundle $V^* := r^*(1)^{-1}\,V$ is said to be the* Bäcklund transform of V of parameters $\alpha, \beta, L^\alpha, L^\beta$. *The q^*-constrained Willmore surface*

$$(\Lambda^{1,0},\, \Lambda^{0,1})^* := (r^*(1)^{-1}\, r^*(\infty)\Lambda^{1,0}, r^*(1)^{-1}\, r^*(0)\Lambda^{0,1})$$

is said to be the Bäcklund transform of $(\Lambda^{1,0}, \Lambda^{0,1})$ of parameters $\alpha, \beta, L^\alpha, L^\beta$.

For further reference, set

$$((\Lambda^*)^{1,0}, (\Lambda^*)^{0,1}) := (r^*(1)^{-1}\, r^*(\infty)\Lambda^{1,0}, r^*(1)^{-1}\, r^*(0)\Lambda^{0,1}).$$

3.3.1. *Bianchi permutability.* Next we establish a *Bianchi permutability* of type p and type q transformations, showing that starting the procedure above with the connections $d^{\lambda,q}_{q_{\beta,L^\beta}}$ (when defined), instead of $d^{\lambda,q}_{p_{\alpha,L^\alpha}}$, produces the same transforms. The underlying argument will play a crucial role when investigating the preservation of reality conditions by Bäcklund transformation, in the next section.

Suppose $\rho L^\beta \cap (L^\beta)^\perp = \{0\}$ and set $\tilde{L}^\alpha_\beta := q_{\beta,L^\beta}(\alpha)L^\alpha$. Suppose, furthermore, that $\rho\tilde{L}^\alpha_\beta \cap (\tilde{L}^\alpha_\beta)^\perp = \{0\}$.[3] Analogously to r^*, we verify that $\hat{r}^* := p_{\alpha,\tilde{L}^\alpha_\beta} q_{\beta,L^\beta}$ satisfies all the hypothesis of Section 3.2 on r. The next result, relating $\hat{r}^*$ to r^*, will be crucial in all that follows.

---

[2] Such a L^β can be obtained by $d^{\beta,q}_V$-parallel transport of $l^\alpha_p \in L^\alpha_p$, with l^α_p non-zero, non-orthogonal to $\rho_p l^\alpha_p$, for some $p \in \Sigma$. Indeed, by (29),

$$(\rho_p p_{\alpha,L^\alpha_p}(\beta) l^\alpha_p, p_{\alpha,L^\alpha_p}(\beta) l^\alpha_p) = (p_{\alpha,L^\alpha_p}(\beta)^{-1}\rho_p l^\alpha_p, p_{\alpha,L^\alpha_p}(\beta) l^\alpha_p) = \frac{(\alpha-\beta)^2}{(\alpha+\beta)^2}(\rho_p l^\alpha_p, l^\alpha_p).$$

[3] This is certainly the case for L^β obtained by $d^{\beta,q}_V$-parallel transport of $l^\alpha_p \in L^\alpha_p$, with l^α_p non-zero, non-orthogonal to $\rho_p l^\alpha_p$, for some $p \in \Sigma$.

Proposition 15.

$$\hat{r}^* = K\, r^*, \tag{32}$$

for $K := q_{\beta,L^\beta}(0)\, q_{\beta,\hat{L}^\beta_\alpha}(0)$.

The proof of the proposition we present next will be based on the following result:

Lemma 16 ([4])**.** *Let* $\gamma(\lambda) = \lambda\, \pi_{L_1} + \pi_{L_0} + \lambda^{-1}\, \pi_{L_{-1}}$ *and* $\hat{\gamma}(\lambda) = \lambda\, \pi_{\hat{L}_1} + \pi_{\hat{L}_0} + \lambda^{-1}\, \pi_{\hat{L}_{-1}}$ *be homomorphisms of* $\mathbb{C}^{n+2}$ *corresponding to decompositions*

$$\mathbb{C}^{n+2} = L_1 \oplus L_0 \oplus L_{-1} = \hat{L}_1 \oplus \hat{L}_0 \oplus \hat{L}_{-1}$$

with $L_{\pm 1}$ *and* $\hat{L}_{\pm 1}$ *null lines and* $L_0 = (L_1 \oplus L_{-1})^\perp$, $\hat{L}_0 = (\hat{L}_1 \oplus \hat{L}_{-1})^\perp$. *Suppose* $\mathrm{Ad}\,\gamma$ *and* $\mathrm{Ad}\,\hat{\gamma}$ *have simple poles. Suppose as well that* ξ *is a map into* $O(\mathbb{C}^{n+2})$ *holomorphic near* 0 *such that* $L_1 = \xi(0)\hat{L}_1$*. Then* $\gamma\xi\hat{\gamma}^{-1}$ *is holomorphic and invertible at* 0.

Now we proceed to the proof of Proposition 15.

Proof. For simplicity, write $p^{-1}_{\mu,L}$ and $q^{-1}_{\mu,L}$ for $\lambda \mapsto p_{\mu,L}(\lambda)^{-1}$ and, respectively, $\lambda \mapsto q_{\mu,L}(\lambda)^{-1}$, in the case $p_{\mu,L}$ and, respectively, $q_{\mu,L}$ are defined. As $L^\alpha = q_{\beta,L^\beta}(\alpha)^{-1}\tilde{L}^\alpha_\beta$, after an appropriate change of variable, we conclude, by Lemma 16, that $p_{\alpha,L^\alpha}\, q^{-1}_{\beta,L^\beta}\, p^{-1}_{\alpha,\tilde{L}^\alpha_\beta}$ admits a holomorphic and invertible extension to $\mathbb{P}^1\backslash\{\pm\beta, -\alpha\}$. On the other hand, in view of (29), the holomorphicity and invertibility of $p_{\alpha,L^\alpha}\, q^{-1}_{\beta,L^\beta}\, p^{-1}_{\alpha,\tilde{L}^\alpha_\beta}$ at the points α and $-\alpha$ are equivalent. Thus $p_{\alpha,L^\alpha}\, q^{-1}_{\beta,L^\beta}\, p^{-1}_{\alpha,\tilde{L}^\alpha_\beta}$ admits a holomorphic and invertible extension to $\mathbb{P}^1\backslash\{\pm\beta\}$, and so does, therefore, $(p_{\alpha,L^\alpha}\, q^{-1}_{\beta,L^\beta}\, p^{-1}_{\alpha,\tilde{L}^\alpha_\beta})^{-1}\, q^{-1}_{\beta,L^\beta}$. A similar argument shows that $p_{\alpha,\tilde{L}^\alpha_\beta}\,(q_{\beta,L^\beta}\, p^{-1}_{\alpha,L^\alpha}\, q^{-1}_{\beta,\hat{L}^\beta_\alpha})$ admits a holomorphic extension to $\mathbb{P}^1\backslash\{\pm\alpha\}$. But $p_{\alpha,\tilde{L}^\alpha_\beta}\, q_{\beta,L^\beta}\, p^{-1}_{\alpha,L^\alpha}\, q^{-1}_{\beta,\hat{L}^\beta_\alpha} = (p_{\alpha,L^\alpha}\, q^{-1}_{\beta,L^\beta}\, p^{-1}_{\alpha,\tilde{L}^\alpha_\beta})^{-1}\, q^{-1}_{\beta,L^\beta}$. We conclude that $p_{\alpha,\tilde{L}^\alpha_\beta}\, q_{\beta,L^\beta}\, p^{-1}_{\alpha,L^\alpha}\, q^{-1}_{\beta,\hat{L}^\beta_\alpha}$ extends holomorphically to $\mathbb{P}^1$ and is, therefore, constant. Evaluating at $\lambda = 0$ gives $p_{\alpha,\tilde{L}^\alpha_\beta}\, q_{\beta,L^\beta}\, p^{-1}_{\alpha,L^\alpha}\, q^{-1}_{\beta,\hat{L}^\beta_\alpha} = q_{\beta,L^\beta}(0)\, q_{\beta,\hat{L}^\beta_\alpha}(0)$, completing the proof. □

According to (30), $\rho\, K\, \rho = K$, showing that K preserves V or, equivalently,

$$K\, V = V. \tag{33}$$

By (32), it follows that

$$r^*(1)^{-1}\, V = \hat{r}^*(1)^{-1}\, V,$$

establishing a *Bianchi permutability* of type p and type q transformations of constrained harmonic bundles, by means of the commutativity of the diagram in Figure 1, below.


V^*
$p_{\alpha,L^\alpha}(1)^{-1}$
$q_{\beta,L^\beta}(1)^{-1}$
$q_{\beta,\hat{L}^\beta_\alpha}(1)^{-1}$
$p_{\alpha,\tilde{L}^\alpha_\beta}(1)^{-1}$
V

FIGURE 1. A Bianchi permutability of type p and type q transformations of constrained harmonic bundles.

Equation (32) makes clear, on the other hand, that

$$\hat{r}^*(1)^{-1}\,\hat{r}^*(\infty)\,\Lambda^{1,0} = r^*(1)^{-1}\,r^*(\infty)\,\Lambda^{1,0}$$

and

$$\hat{r}^*(1)^{-1}\,\hat{r}^*(0)\,\Lambda^{0,1} = r^*(1)^{-1}\,r^*(0)\,\Lambda^{0,1}.$$

We conclude that, despite not coinciding, r^* and $\hat{r}^*$ produce the same transforms of (constrained) harmonic bundles and Willmore surfaces. As a final remark, note that, yet again by equation (32),

$$\hat{q}^* := \mathrm{Ad}_{\hat{r}^*(1)^{-1}}(\mathrm{Ad}_{\hat{r}^*(0)}q^{1,0} + \mathrm{Ad}_{\hat{r}^*(\infty)}q^{0,1}) = q^*.$$

3.3.2. *Real Bäcklund transformation.* As we verify next, Bäcklund transformation preserves reality conditions, for special choices of parameters.

Suppose V is a real q-constrained harmonic bundle. Obviously, the reality of V establishes that of ρ and, therefore,

$$\overline{p_{\mu,L}(\lambda)} = p_{\overline{\mu},\overline{L}}\,(\overline{\lambda}), \qquad \overline{q_{\mu,L}(\lambda)} = q_{\overline{\mu},\overline{L}}\,(\overline{\lambda}), \tag{34}$$

for all μ, L and $\lambda \in \mathbb{C}\backslash\{\pm\mu\}$.

Lemma 17. *Suppose $\alpha \in \mathbb{C}\backslash(S^1 \cup \{0\})$. Then we can choose $\beta = \overline{\alpha}^{\,-1}$ and $L^\beta = \overline{L^\alpha}$ and both r^* and $\hat{r}^*$ are defined.*

Before proceeding to the proof of the lemma, note that

Proof. The reality of ρ makes it clear that the non-orthogonality of L^α and ρL^α establishes that of $\overline{L^\alpha}$ and $\rho\overline{L^\alpha}$, as well as, together with (34) and (28), that, if $\rho p_{\alpha,L^\alpha}(\overline{\alpha}^{\,-1})\overline{L^\alpha} \cap p_{\alpha,L^\alpha}(\overline{\alpha}^{\,-1})\overline{L^\alpha} = \{0\}$, then $\rho q_{\overline{\alpha}^{\,-1},\overline{L^\alpha}}(\alpha)L^\alpha \cap q_{\overline{\alpha}^{\,-1},\overline{L^\alpha}}(\alpha)L^\alpha = \{0\}$. On the other hand, the reality of V establishes that of $\mathcal{D}_V$ and $\mathcal{N}_V$, so that, by the reality of q, $d_V^{\overline{\alpha}^{\,-1},q} = \overline{d_V^{\alpha,q}}$. Hence the $d_V^{\alpha,q}$-parallelness of L^α establishes the $d_V^{\overline{\alpha}^{\,-1},q}$-parallelness of $\overline{L^\alpha}$. Obviously, if α is non-unit, then $\overline{\alpha}^{\,-1} \neq \pm\alpha$. We are left to verify that we can choose L^α a $d_V^{\alpha,q}$-parallel null line subbundle of $\underline{\mathbb{C}}^{n+2}$ such that, locally, $\rho L^\alpha \cap L^\alpha = \{0\}$ and $\rho p_{\alpha,L^\alpha}(\overline{\alpha}^{\,-1})\overline{L^\alpha} \cap p_{\alpha,L^\alpha}(\overline{\alpha}^{\,-1})\overline{L^\alpha} = \{0\}$. For this, let v and w be sections of V and $V^\perp$, respectively, with (v,v) never-zero, $(v,\overline{v}) = 0$ and $(w,w) = -(v,v)$.

Define a null section of $\underline{\mathbb{C}}^{n+2}$ by $l^\alpha := v + w$ and then $L^\alpha \subset \underline{\mathbb{C}}^{n+2}$ by $d_V^{\alpha,q}$-parallel transport of l^α_p, for some point $p \in \Sigma$. $\square$

Let us focus then on the particular case of Bäcklund transformation of parameters $\alpha, \beta, L^\alpha, L^\beta$ with

$$\alpha \in \mathbb{C}\backslash(S^1 \cup \{0\}),\ \ \beta = \overline{\alpha}^{\,-1},\ \ L^\beta = \overline{L^\alpha},$$

which we refer to as *Bäcklund transformation of parameters* α, L^α. For this particular choice of parameters, we write $\tilde{L}^\alpha$ and $\hat{L}_\alpha$ for $\tilde{L}^\alpha_\beta$ and $\hat{L}^\beta_\alpha$, respectively. Note that, by (28) and (34), $\overline{\hat{L}_\alpha} = \tilde{L}^\alpha$. On the other hand,

$$\overline{r^*(1)^{-1}} = \overline{p_{\alpha,L^\alpha}(1)^{-1}}\,\overline{q_{\beta,\hat{L}_\alpha}(1)^{-1}} = q_{\beta,L^\beta}(1)^{-1}p_{\alpha,\overline{\hat{L}_\alpha}}(1)^{-1},$$

whilst, by (32),

$$r^*(1)^{-1} = (K^{-1}\hat{r}^*(1))^{-1} = q_{\beta,L^\beta}(1)^{-1}p_{\alpha,\tilde{L}^\alpha}(1)^{-1}K.$$

Hence

$$\overline{r^*(1)^{-1}} = r^*(1)^{-1}K^{-1}. \tag{35}$$

By (33), it follows that $\overline{V^*} = V^*$. Next we establish the reality of q^*. Yet again by (34), $\overline{r^*(0)} = q_{\alpha^{-1},\overline{\hat{L}_\alpha}}(0) = p_{\alpha^{-1},\overline{\hat{L}_\alpha}}(\infty)$ and, on the other hand, by (32), $r^*(\infty) = K^{-1}p_{\alpha,\tilde{L}_\alpha}(\infty)$, so that

$$\overline{r^*(0)} = Kr^*(\infty). \tag{36}$$

Together with (35), this makes clear that $\overline{(q^*)^{1,0}} = (q^*)^{0,1}$, the reality of q establishes that of q^*.

We conclude that:

Theorem 18. *If V is a real q-constrained harmonic bundle, then the Bäcklund transform V^* of V, of parameters α, L^α, is a real q^*-constrained harmonic bundle.*

A real transformation on the level of constrained Willmore surface follows. Equation (32) plays, yet again, a crucial role, by showing that

$$r^*(1) = K^{-1}q_{\alpha^{-1},\tilde{L}^\alpha}(1)p_{\overline{\alpha},L^\beta}(1) = K^{-1}\overline{q_{\beta,\hat{L}_\alpha}(1)p_{\alpha,L^\alpha}(1)} = K^{-1}\overline{r^*(1)},$$

and, therefore,

$$\overline{r^*(1)^{-1}} = r^*(1)^{-1}\overline{K}. \tag{37}$$

Suppose $(\Lambda^{1,0}, \Lambda^{0,1})$ is a real surface, so that, in particular, $\overline{\Lambda^{1,0}} = \Lambda^{0,1}$. By (36) and (37), it follows that $\overline{(\Lambda^*)^{1,0}} = (\Lambda^*)^{0,1}$, establishing the reality of the bundle

$$\Lambda^* := (\Lambda^*)^{1,0} \cap (\Lambda^*)^{0,1}.$$

We conclude that:

Theorem 19. *If Λ is a real q-constrained Willmore surface, then the Bäcklund transform Λ^* of Λ, of parameters α, L^α, is a real q^*-constrained Willmore surface.*

3.4. **Spectral deformation vs Bäcklund transformation.** Bäcklund transformation and spectral deformation permute, as follows:

Theorem 20. *Let* α,β,L^α,L^β *be Bäcklund transformation parameters to* V*,* $\lambda\in\mathbb{C}\backslash\{0,\pm\alpha,\pm\beta\}$ *and* $\phi^\lambda:(\underline{\mathbb{C}}^{n+2},d_V^{\lambda;q})\rightarrow(\underline{\mathbb{C}}^{n+2},d)$ *be an isometry of bundles preserving connections. The Bäcklund transform of parameters* $\frac{\alpha}{\lambda},\frac{\beta}{\lambda}$*,* $\phi^\lambda L^\alpha$*,* $\phi^\lambda L^\beta$ *of the spectral deformation* $\phi^\lambda V$ *of* V*, of parameter* λ*, corresponding to the multiplier* q*, coincides with the spectral deformation of parameter* λ*, corresponding to the multiplier* q^**, of the Bäcklund transform of parameters* $\alpha,\beta,L^\alpha,L^\beta$ *of* V*. Furthermore, if* $\phi^\lambda_*:(\underline{\mathbb{C}}^{n+2},d_{V^*}^{\lambda;q^*})\rightarrow(\underline{\mathbb{C}}^{n+2},d)$ *is an isometry preserving connections, then the diagram in Figure 2 commutes.*

$$\begin{tikzcd}
& (\phi^\lambda_*(\Delta^*)^{1,0},\phi^\lambda_*(\Delta^*)^{0,1})=(\phi^\lambda\Delta^{1,0},\phi^\lambda\Delta^{0,1})^* \arrow[dl, dash, "\lambda"'] \arrow[dr, dash, "{\frac{\alpha}{\lambda},\frac{\beta}{\lambda},\phi^\lambda L^\alpha,\phi^\lambda L^\beta}"] & \\
(\Delta^{1,0},\Delta^{0,1})^* \arrow[dr, dash, "{\alpha,\beta,L^\alpha,L^\beta}"'] & & (\phi^\lambda\Delta^{1,0},\phi^\lambda\Delta^{0,1}) \arrow[dl, dash, "\lambda"] \\
& (\Delta^{1,0},\Delta^{0,1}) &
\end{tikzcd}$$

FIGURE 2. A Bianchi permutability of spectral deformation and Bäcklund transformation of constrained Willmore surfaces.

Proof. It is trivial, noting that $\phi^\lambda r^*(\lambda)^{-1}r^*(1):(\underline{\mathbb{C}}^{n+2},d_{V^*}^{\lambda;q^*})\rightarrow(\underline{\mathbb{C}}^{n+2},d)$ is an isometry of bundles preserving connections. $\square$

For $\lambda\in\{\pm\alpha,\pm\beta\}$, it is not clear how the spectral deformation of parameter λ relates to the Bäcklund transformation of parameters $\alpha,\beta,L^\alpha,L^\beta$.

3.5. **Isothermic condition under constrained Willmore transformation.** Equation (1) makes it clear that, although the second fundamental form is not conformally invariant, conformal curvature line coordinates are preserved under conformal changes of the metric and, therefore, so is the isothermic surface condition. The next result presents a manifestly conformally-invariant formulation of the isothermic surface condition, established in [6] and discussed also in [11].

Lemma 21 ([6],[11])**.** Λ *is isothermic if and only if there exists a non-zero closed real* 1*-form* $\eta\in\Omega^1(\Lambda\wedge\Lambda^{(1)})$*. Under these conditions, we may say that* (Λ,η) *is an isothermic surface. The form* η *is defined up to a real constant scale.*

Remark 3. *According to the decomposition* (12)*, given* $\eta \in \Omega^1(\Lambda \wedge \Lambda^{(1)})$*,* $d\eta = d^{\mathcal{D}}\eta + [\mathcal{N} \wedge \eta]$ *vanishes if and only if* $d^{\mathcal{D}}\eta = 0 = [\mathcal{N} \wedge \eta]$*.*

Proposition 22. *A constrained Willmore surface has a unique multiplier if and only if it is not an isothermic surface. Furthermore:*

i) if $q_1 \neq q_2$ *are multipliers for* Λ*, then* $(\Lambda, *(q_1 - q_2))$ *is isothermic;*

ii) if (Λ, η) *is an isothermic* q*-constrained Willmore surface, then the set of multipliers to* Λ *is the affine space* $q + \langle *\eta \rangle_{\mathbb{R}}$*.*

Proof. It is immediate, noting that $[\mathcal{N} \wedge \eta] = [*\eta \wedge *\mathcal{N}]$ and recalling Proposition 4 - *iii*). ☐

The isothermic surface condition is known to be preserved under constrained Willmore spectral deformation:

Proposition 23 ([8])**.** *Constrained Willmore spectral deformation preserves the isothermic surface condition.*

Next we derive it in our setting.

Proof. Suppose (Λ, η) is an isothermic q-constrained Willmore surface, for some $\eta, q \in \Omega^1(\Lambda \wedge \Lambda^{(1)})$. Fix $\lambda \in S^1$ and $\phi^\lambda_q : (\underline{\mathbb{R}}^{n+1,1}, d^\lambda_q) \to (\underline{\mathbb{R}}^{n+1,1}, d)$ an isometry preserving connections. Set

$$\eta_\lambda := \lambda^{-1}\eta^{1,0} + \lambda\, \eta^{0,1}.$$

To prove the theorem, we show that $(\phi^\lambda_q\Lambda, \mathrm{Ad}_{\phi^\lambda_q}\eta_\lambda)$ is isothermic. Obviously, the reality of η establishes that of $\mathrm{Ad}_{\phi^\lambda_q}\eta_\lambda$. Recall (15) to conclude that $(\phi^\lambda_q\Lambda)^{(1)} = \phi^\lambda_q\Lambda^{(1)}$ and, therefore, that $\mathrm{Ad}_{\phi^\lambda_q}\eta_\lambda$ takes values in $\phi^\lambda_q\Lambda \wedge (\phi^\lambda_q\Lambda)^{(1)}$. Now observe that, given $a, b \in \mathbb{R}^{n+1,1}$ and $T \in o(\mathbb{R}^{n+1,1})$, $[T, a \wedge b] = (Ta) \wedge b + a \wedge (Tb)$, to conclude that $[\Lambda \wedge \Lambda^{(1)}, \Lambda \wedge \Lambda^{(1)}] \subset \Lambda \wedge \Lambda = \{0\}$. Thus $[q^{1,0} \wedge \eta^{0,1}] = 0 = [q^{0,1} \wedge \eta^{1,0}]$ and, therefore,

$$\begin{aligned} d^{d^\lambda_q}\eta_\lambda &= d^{\mathcal{D}}\eta_\lambda + [(\lambda^{-1}\mathcal{N}^{1,0} + \lambda\mathcal{N}^{0,1} + (\lambda^{-2} - 1)q^{1,0} + (\lambda^2 - 1)q^{0,1}) \wedge \eta_\lambda] \\ &= d^{\mathcal{D}}\eta_\lambda + [\mathcal{N} \wedge \eta]. \end{aligned}$$

According to the decomposition (12), we conclude that $d(\mathrm{Ad}_{\phi^\lambda_q}\eta_\lambda) = \phi^\lambda_q \circ d^{d^\lambda_q}\eta_\lambda \circ (\phi^\lambda_q)^{-1}$ vanishes if and only if $d^{\mathcal{D}}\eta_\lambda = 0 = [\mathcal{N} \wedge \eta]$. Remark 3 and Lemma 4 complete the proof. ☐

As for Bäcklund transformation of constrained Willmore surfaces, we believe it does not necessarily preserve the isothermic condition. This shall be the subject of further work. In contrast, the constancy of the mean curvature of a surface in 3-dimensional space-form is preserved by both constrained Willmore spectral deformation and Bäcklund transformation, for special choices of parameters, with preservation of both the space-form and the mean curvature in the latter case, cf. [16].

References

[1] W. Blaschke, *Vorlesungen über Differentialgeometrie III: Differentialgeometrie der Kreise und Kugeln*, Grundlehren XXIX, Springer, Berlin (1929).

[2] C. Bohle, G. P. Peters and U. Pinkall, *Constrained Willmore Surfaces*, Calculus of Variations and Partial Differential Equations **32** (2008) 263-277.

[3] R. Bryant, *A duality theorem for Willmore surfaces*, Journal of Differential Geometry 20 (1984) 23-53.

[4] F. E. Burstall, *Isothermic surfaces: Conformal geometry, Clifford algebras and Integrable systems*, Integrable systems, Geometry and Topology (ed. C.-L. Terng), Volume **36**, American Mathematical Society/International Press Studies in Advanced Mathematics, American Mathematical Society, Providence (2006) 1-82.

[5] F. E. Burstall and D. M. J. Calderbank, *Conformal Submanifold Geometry I-III*, arXiv:1006.5700v1 [math.DG] (2010).

[6] F. E. Burstall, N. M. Donaldson, F. Pedit, U. Pinkall, *Isothermic Submanifolds of Symmetric R-Spaces*, J. reine u. Angew. Math. (to appear). DOI: 10.1515/CRELLE.2011.075.

[7] F. E. Burstall, D. Ferus, K. Leschke, F. Pedit, and U. Pinkall, *Conformal Geometry of Surfaces in* S^4 *and Quaternions*, Lecture Notes in Mathematics, vol. 1772, Springer Verlag, Heidelberg, Berlin, New York, 2002.

[8] F. E. Burstall, F. Pedit and U. Pinkall, *Schwarzian Derivatives and Flows of Surfaces*, Contemporary Mathematics **308** (2002) 39-61.

[9] F. E. Burstall and A. C. Quintino, *Dressing Transformations of Constrained Willmore Surfaces* (in preparation).

[10] F. E. Burstall and J. Rawnsley, *Twistor theory for Riemannian symmetric spaces*, Lecture Notes in Mathematics 1424, Springer-Verlag (1990).

[11] F. E. Burstall and S. D. Santos, *Special Isothermic Surfaces of Type* d, Journal of the London Mathematical Society (to appear). arXiv:1006.3175v1 [math.DG].

[12] J. Darboux, *Leçons sur la Théorie Générale des Surfaces et les Applications Géometriques du Calcul Infinitésimal, Parts 1 and 2*, Gauthier-Villars, Paris, 1887.

[13] N. Ejiri, *Willmore Surfaces with a Duality in* $S^n(1)$, Proceedings of the London Mathematical Society (3) **57** (1988) 383-416.

[14] S. Germain, *Recherches sur la théorie des surfaces élastiques*, Courcier, Paris (1821).

[15] S. Germain, *Remarques sur la nature, les bornes et l'etendue de la question des surfaces élastiques, et equation generale de ces surfaces*, Courcier, Paris (1826).

[16] A. C. Quintino, *Constrained Willmore Surfaces: Symmetries of a Möbius Invariant Integrable System* - Based on the author's PhD Thesis. arXiv:0912.5402v1 [math.DG] (2009).

[17] J. Richter, *Conformal Maps of a Riemann Surface into the Space of Quaternions*, PhD thesis, Technischen Universität Berlin (1997).

[18] M. Rigoli, *The Conformal Gauss Map of Submanifolds of the Möbius Space*, Annals of Global Analysis and Geometry **5**, N?. 2 (1987) 97-116.

[19] C.-L. Terng and K. Uhlenbeck, *Bäcklund Transformations and Loop Group Actions*, Communications on Pure and Applied Mathematics **53** (2000) 1-75.

[20] G. Thomsen, Ueber konforme Geometrie I, *Grundlagen der Konformen Flaechentheorie*, Abhandlungen aus dem Mathematischen Seminar der Universit?t Hamburg **3** (1923) 31-56.

[21] K. Uhlenbeck, *Harmonic Maps into Lie Groups (Classical Solutons of the Chiral Model)*, Journal of Differential Geometry **30** (1989) 1-50.

[22] T. Willmore, *Note on Embedded Surfaces*, Analele Stiintifice ale Universitatii "Alexandru Ioan Cuza" din Iasi, N. Ser., Sect. Ia **11**B (1965) 493-496.

[23] T. Willmore, *Riemannian Geometry*, Oxford Science Publications (1993).

A. C. Quintino, Centro de Matemática e Aplicações Fundamentais, Universidade de Lisboa, 1649-003 Lisboa, Portugal, *e-mail address:* aurea@ptmat.fc.ul.pt